

ON WEAKLY 1-ABSORBING PRIMARY IDEALS OF COMMUTATIVE RINGS

AYMAN BADAWI AND ECE YETKIN CELIKEL

ABSTRACT. Let R be a commutative ring with $1 \neq 0$. In this paper, we introduce the concept of weakly 1-absorbing primary ideal which is a generalization of 1-absorbing primary ideal. A proper ideal I of R is called a *weakly 1-absorbing primary* ideal if whenever nonunit elements $a, b, c \in R$ and $0 \neq abc \in I$, then $ab \in I$ or $c \in \sqrt{I}$. A number of results concerning weakly 1-absorbing primary ideals and examples of weakly 1-absorbing primary ideals are given. Furthermore, we give the correct version of a result on 1-absorbing primary ideals of commutative rings.

1. INTRODUCTION

Throughout this paper, all rings are commutative with nonzero identity. Let R be a commutative ring. By a proper ideal I of R , we mean an ideal I of R with $I \neq R$. Let I be a proper ideal of R . By \sqrt{I} , we mean the *radical* of I , that is, $\{a \in R \mid a^n \in I \text{ for some positive integer } n\}$. In particular, $\sqrt{\{0\}}$ denotes the set of all nilpotent elements of R . We define $Z_I(R) = \{r \in R \mid rs \in I \text{ for some } s \in R \setminus I\}$. A ring R is called a *reduced* ring if it has no nonzero nilpotent elements; i.e., $\sqrt{\{0\}} = \{0\}$. For two ideals I and J of R , the *residual division* of I and J is defined to be the ideal $(I : J) = \{a \in R \mid aJ \subseteq I\}$. Let R be a commutative ring with identity and M a unitary R -module. Then $R(+M) = R \times M$ with coordinate-wise addition and multiplication $(a, m)(b, n) = (ab, an + bm)$ is a commutative ring with identity $(1, 0)$ called the *idealization* of M . A ring R is called a *quasilocal* ring if R has exactly one maximal ideal. As usual, \mathbb{Z} and \mathbb{Z}_n will denote the ring of integers and integers modulo n , respectively.

Since prime and primary ideals have key roles in commutative ring theory, many authors have studied generalizations of prime and primary ideals (see [1], [2], [3], [5], [6],[7], [8], [9], [10], and [11]). Anderson and Smith introduced in [2] the notion of weakly prime ideals. A proper ideal I of R is called a *weakly prime* ideal of R if whenever $a, b \in R$ and $0 \neq ab \in I$, then $a \in I$ or $b \in I$. Then Atani and Farzalipour introduced in [5] the concept of weakly primary ideals. A proper ideal I of R is called a *weakly primary* ideal of R if whenever $a, b \in R$ and $0 \neq ab \in I$, then $a \in I$ or $b \in \sqrt{I}$. For a different generalization of prime ideals and weakly prime ideals, the contexts of 2-absorbing and weakly 2-absorbing ideals were defined. According to [6] and [7], a proper ideal I of R is called a *2-absorbing (weakly 2-absorbing)* ideal of a commutative ring R , if whenever $a, b, c \in R$ and $abc \in I$ ($0 \neq abc \in I$),

2000 *Mathematics Subject Classification.* Primary 13A15, 13F05; Secondary 05A15, 13G05.

Key words and phrases. 1-absorbing primary ideal, 2-absorbing primary ideal, 2-absorbing ideal, weakly 2-absorbing primary ideal, weakly primary.

then $ab \in I$ or $bc \in I$ or $ac \in I$. As a generalization of 2-absorbing and weakly 2-absorbing ideals, 2-absorbing primary and weakly 2-absorbing primary ideals were defined in [8] and [9], respectively. A proper ideal I of a commutative ring R is said to be *2-absorbing primary* (*weakly 2-absorbing primary*) if whenever $a, b, c \in R$ and $abc \in I$ ($0 \neq abc \in I$), then $ab \in I$ or $bc \in \sqrt{I}$ or $ac \in \sqrt{I}$. In a recent study [10], we call a proper ideal I of a commutative ring R a *1-absorbing primary* ideal if whenever nonunit elements $a, b, c \in R$ and $abc \in I$, then $ab \in I$ or $c \in \sqrt{I}$.

In this paper, we introduce the concept of weakly 1-absorbing primary ideal of a commutative ring R . A proper ideal I of a commutative ring R is called a *weakly 1-absorbing primary* ideal of R if whenever nonunit elements $a, b, c \in R$ and $0 \neq abc \in I$, then $ab \in I$ or $c \in \sqrt{I}$. It is clear that a 1-absorbing primary ideal of a commutative ring R is a weakly 1-absorbing primary ideal of R . However, since $\{0\}$ is always weakly 1-absorbing primary, a weakly 1-absorbing primary ideal of a commutative ring R needs not be a 1-absorbing primary ideal of R (see Example 1).

Among many results, we show (Theorem 2) that if a proper ideal I of a commutative ring R is a weakly 1-absorbing primary ideal of R such that \sqrt{I} is a maximal ideal of R , then I is a primary ideal of R , and hence I is a 1-absorbing primary ideal of R . We show (Theorem 3) that if R is a commutative reduced ring and I is a weakly 1-absorbing primary ideal of R , then \sqrt{I} is a prime ideal of R . If I is a proper nonzero ideal of a commutative von Neumann regular ring R , then we show (Theorem 4) that I is a weakly 1-absorbing primary ideal of R if and only if I is a 1-absorbing primary ideal of R , if and only if I is a primary ideal of R . We show (Theorem 5) that if R be a commutative non-quasilocal ring and I is a proper ideal of R such that $\text{ann}(i) = \{r \in R \mid ri = 0\}$ is not a maximal ideal of R for every element $i \in I$, then I is a weakly 1-absorbing primary ideal of R if and only if I is a weakly primary ideal of R . If I is a proper ideal of a commutative reduced divided ring R , then we show (Theorem 7) that I is a weakly 1-absorbing primary ideal of R if and only if I is a weakly primary ideal of R . If I is a weakly 1-absorbing primary ideal of a commutative ring R that is not a 1-absorbing primary ideal of R , then we give (Theorem 10) sufficient conditions so that $I^3 = \{0\}$ (i.e., $I \subseteq \sqrt{\{0\}}$). In Theorem 9, we obtain some equivalent conditions for weakly 1-absorbing primary ideals of u -rings. We give (Theorem 13) a characterization of weakly 1-absorbing primary ideals in $R = R_1 \times R_2$, where R_1 and R_2 are commutative rings with identity that are not fields. If R_1, R_2, \dots, R_n are commutative rings with identity for some $2 \leq n < \infty$ and $R = R_1 \times \dots \times R_n$, then it is shown (Theorem 14) that every proper ideal of R is a weakly 1-absorbing primary ideal of R if and only if $n = 2$, and R_1, R_2 are fields. For a weakly 1-absorbing primary ideal of a ring R , we show (Theorem 17) that $S^{-1}I$ is a weakly 1-absorbing primary ideal of $S^{-1}R$ for every multiplicatively closed subset S of R that is disjoint from I , and we show that the converse holds if $S \cap Z(R) = S \cap Z_I(R) = \emptyset$. We give (Remark 1) the correct versions of [10, Theorem 17(1), Corollary 3, and Corollary 4].

2. PROPERTIES OF WEAKLY 1-ABSORBING PRIMARY IDEALS

Definition 1. *Let R be a commutative ring and I be a proper ideal of R . We call I a weakly 1-absorbing primary ideal of R if whenever nonunit elements $a, b, c \in R$ and $0 \neq abc \in I$, then $ab \in I$ or $c \in \sqrt{I}$.*

It is clear that every 1-absorbing primary ideal of a commutative ring R is a weakly 1-absorbing primary ideal of R , and $I = \{0\}$ is a weakly 1-absorbing primary ideal of R . In the following example, we construct a weakly 1-absorbing primary ideal of a commutative ring R that is neither 1-absorbing primary nor weakly primary.

- Example 1.** (1) $I = \{0\}$ is a weakly 1-absorbing primary ideal of $R = \mathbb{Z}_6$ that is not a 1-absorbing primary ideal of R . Indeed, $2 \cdot 2 \cdot 3 \in I$, but neither $2 \cdot 2 \in I$ nor $3 \in \sqrt{I}$. Note that I is a weakly primary ideal of R .
- (2) Let $A = \mathbb{Z}_2[[X, Y]]$, $I = (XY^2, YX^2)A$, $R = A/I$, and $J = (XY)A/I$. We show that J is a weakly 1-absorbing primary ideal of R that is neither 1-absorbing primary nor weakly primary. Assume that $abc \in J$ for some nonunit elements $a, b, c \in R$. Then $abc = XYZ + I$ for some nonunit element $Z \in A$. Hence $abc = I \in J$ by construction of J . Thus J is a weakly 1-absorbing primary ideal of R . Since $(X+I)(X+I)(Y+I) = I \in J$ and neither $X^2 + I \in J$ nor $Y + I \in \sqrt{J}$, we conclude that J is not a 1-absorbing primary ideal of R . Since $I \neq (X+I)(Y+I) = XY + I \in J$ and neither $X+I \in J$ nor $Y+I \in \sqrt{J}$, we conclude that J is not a weakly primary ideal of R .

We begin with the following trivial result without proof.

Theorem 1. Let I be a proper ideal of a commutative ring R .

- (1) If I is a weakly prime ideal, then I is a weakly 1-absorbing primary ideal.
- (2) If I is a weakly primary ideal, then I is a weakly 1-absorbing primary ideal.
- (3) If I is a 1-absorbing primary ideal, then I is a weakly 1-absorbing primary ideal.
- (4) If I is a weakly 1-absorbing primary ideal, then I is a weakly 2-absorbing primary ideal.
- (5) If R is an integral domain, then I is a weakly 1-absorbing primary ideal if and only if I is a 1-absorbing primary ideal of R .
- (6) Let R be a quasilocal ring with maximal ideal $\sqrt{\{0\}}$. Then every proper ideal of R is a weakly 1-absorbing primary ideal of R .

We recall that a proper ideal I of a commutative ring R is called a *semiprimary* ideal of R if \sqrt{I} is a prime ideal of R . For an interesting article on semiprimary ideals of commutative rings, see [12]. For a recent related article on semiprimary ideals, we recommend [11]. We have the following result.

Theorem 2. Let R be a commutative ring and I be a weakly 1-absorbing primary ideal of R . If \sqrt{I} is a maximal ideal of R , then I is a primary ideal of R , and hence I is a 1-absorbing primary ideal of R . In particular, if I is a weakly 1-absorbing primary ideal of R that is not a 1-absorbing primary ideal of R , then \sqrt{I} is not a maximal ideal of R .

Proof. Suppose that \sqrt{I} is a maximal ideal of R . Then I is a semiprimary ideal of R . Since I is a semiprimary ideal of R and \sqrt{I} is a maximal ideal of R , we conclude that I is a primary ideal of R by [14, p. 153]. Thus I is a 1-absorbing primary ideal of R . \square

Theorem 3. Let R be a commutative reduced ring. If I is a nonzero weakly 1-absorbing primary ideal of R , then \sqrt{I} is a prime ideal of R . In particular, if \sqrt{I} is

a maximal ideal of R , then I is a primary ideal of R , and hence I is a 1-absorbing primary ideal of R .

Proof. Suppose that $0 \neq ab \in \sqrt{I}$ for some $a, b \in R$. We may assume that a, b are nonunits. Then there exists an even positive integer $n = 2m$ ($m \geq 1$) such that $(ab)^n \in I$. Since $\sqrt{\{0\}} = \{0\}$, we have $(ab)^n \neq 0$. Hence $0 \neq a^m a^m b^n \in I$. Thus $a^m a^m = a^n \in I$ or $b^n \in \sqrt{I}$, and therefore \sqrt{I} is a weakly prime ideal of R . Since R is reduced and $I \neq \{0\}$, we conclude that \sqrt{I} is a prime ideal of R by [2, Corollary 2]. The proof of the ‘‘in particular’’ statement is now clear by Theorem 2. \square

Recall that a commutative ring R is called a *von Neumann regular* ring if and only if for every $x \in R$, there is a $y \in R$ such that $x^2y = x$. It is known that a commutative ring R is a von Neumann regular ring if and only if for each $x \in R$, there is an idempotent $e \in R$ and a unit $u \in R$ such that $x = eu$. For a recent article on von Neumann regular rings, see[4]. We have the following result.

Theorem 4. *Let R be a commutative von Neumann regular ring and I be a nonzero ideal of R . Then the following statements are equivalent.*

- (1) I is a weakly 1-absorbing primary ideal of R .
- (2) I is a primary ideal of R .
- (3) I is a 1-absorbing primary ideal of R .

Proof. (1) \Rightarrow (2) Since R is a commutative von Neumann regular ring, we know that R is reduced. Hence \sqrt{I} is a prime ideal of R by Theorem 3. Since every prime ideal of a von Neumann regular ring is maximal, we conclude that \sqrt{I} is a maximal ideal of R . Hence I is a primary ideal of R by Theorem 2.

(2) \Rightarrow (3) \Rightarrow (1) It is clear. \square

Let A, I, R , and J be as in Example 1(2). Then R is a quasilocal ring with maximal ideal $M = (X, Y)A/I$, and $\text{ann}(XY+I) = \{a \in R \mid a(XY+I) = 0\} = M$. We have the following result.

Theorem 5. *Let R be a commutative non-quasilocal ring and I be a proper ideal of R such that $\text{ann}(i) = \{r \in R \mid ri = 0\}$ is not a maximal ideal of R for every element $i \in I$. Then I is a weakly 1-absorbing primary ideal of R if and only if I is a weakly primary ideal of R .*

Proof. If I is a weakly primary ideal of R , then I is a weakly 1-absorbing primary ideal of R by Theorem 1(2). Hence suppose that I is a weakly 1-absorbing primary ideal of R and suppose that $0 \neq ab \in I$ for some elements $a, b \in R$. We show that $a \in I$ or $b \in \sqrt{I}$. We may assume that a, b are nonunit elements of R . Let $\text{ann}(ab) = \{c \in R \mid cab = 0\}$. Since $ab \neq 0$, $\text{ann}(ab)$ is a proper ideal of R . Let L be a maximal ideal of R such that $\text{ann}(ab) \subsetneq L$. Since R is a non-quasilocal ring, there is a maximal ideal M of R such that $M \neq L$. Let $m \in M \setminus L$. Hence $m \notin \text{ann}(ab)$ and $0 \neq mab \in I$. Since I is a weakly 1-absorbing primary ideal of R , we have $ma \in I$ or $b \in \sqrt{I}$. If $b \in \sqrt{I}$, then we are done. Hence assume that $b \notin \sqrt{I}$. Hence $ma \in I$. Since $m \notin L$ and L is a maximal ideal of R , we conclude that $m \notin J(R)$. Hence there exists an $r \in R$ such that $1 + rm$ is a nonunit element of R . Suppose that $1 + rm \notin \text{ann}(ab)$. Hence $0 \neq (1 + rm)ab \in I$. Since I is a weakly 1-absorbing primary ideal of R and $b \notin \sqrt{I}$, we conclude that $(1 + rm)a = a + rma \in I$. Since $rma \in I$, we have $a \in I$ and we are done. Suppose that $1 + rm \in \text{ann}(ab)$. Since $\text{ann}(ab)$ is not a maximal ideal of R and $\text{ann}(ab) \subsetneq L$, there is a $w \in L \setminus \text{ann}(ab)$.

Hence $0 \neq wab \in I$. Since I is a weakly 1-absorbing primary ideal of R and $b \notin \sqrt{I}$, we conclude that $wa \in I$. Since $1 + rm \in \text{ann}(ab) \subsetneq L$ and $w \in L \setminus \text{ann}(ab)$, we have $1 + rm + w$ is a nonzero nonunit element of L . Hence $0 \neq (1 + rm + w)ab \in I$. Since I is a weakly 1-absorbing primary ideal of R and $b \notin \sqrt{I}$, we conclude that $(1 + rm + w)a = a + rma + wa \in I$. Since $rma, wa \in I$, we conclude that $a \in I$. \square

Question. Is Theorem 5 still valid without the assumption that $\text{ann}(i) = \{r \in R \mid ri = 0\}$ is not a maximal ideal of R for every element $i \in I$? We are unable to give a proof of Theorem 5 without this assumption.

In light of the proof of Theorem 5, we have the following result.

Theorem 6. *Let I be a weakly 1-absorbing primary ideal of a commutative ring R such that for every nonzero element $i \in I$, there exists a nonunit $w \in R$ such that $wi \neq 0$ and $w + u$ is a nonunit element of R for some unit $u \in R$. Then I is a weakly primary ideal of R .*

Proof. Suppose that $0 \neq ab \in I$ and $b \notin \sqrt{I}$ for some $a, b \in R$. We may assume that a, b are nonunit elements of R . Hence there is a nonunit $w \in R$ such that $wab \neq 0$ and $w + u$ is a nonunit element of R for some unit $u \in R$. Since $0 \neq wab \in I$, $b \notin \sqrt{I}$, and I is a weakly 1-absorbing primary ideal of R , we conclude that $wa \in I$. Since $0 \neq (w + u)ab \in I$, I is a weakly 1-absorbing primary ideal of R , and $b \notin \sqrt{I}$, we conclude that $wa + ua = (w + u)a \in I$. Since $wa \in I$ and $wa + ua \in I$, we conclude that $ua \in I$. Since u is a unit, we have $a \in I$. \square

Corollary 1. *Let R be a commutative ring and $A = R[X]$. Suppose that I is a weakly 1-absorbing primary ideal of A . Then I is a weakly primary ideal of A .*

Proof. Since $Xi \neq 0$ for every nonzero $i \in I$ and $X + 1$ is a nonunit element of A , we are done by Theorem 6. \square

Recall that a commutative ring R is called *divided* if for every prime ideal P of R and for every $x \in R \setminus P$, we have $x \mid p$ for every $p \in P$. We have the following result.

Theorem 7. *Let R be a commutative reduced divided ring and I be a proper ideal of R . Then the following statements are equivalent.*

- (1) I is a weakly 1-absorbing primary ideal of R .
- (2) I is a weakly primary ideal of R .

Proof. (1) \Rightarrow (2) Suppose that $0 \neq ab \in I$ for some $a, b \in R$ and $b \notin \sqrt{I}$. We may assume that a, b are nonunit elements of R . Since \sqrt{I} is a prime ideal of R by Theorem 3, we conclude that $a \in \sqrt{I}$. Since R is divided, we conclude that $b \mid a$. Thus $a = bc$ for some $c \in R$. Observe that c is a nonunit element of R as $b \notin \sqrt{I}$ and $a \in \sqrt{I}$. Since $0 \neq ab = bcb \in I$, I is weakly 1-absorbing primary, and $b \notin \sqrt{I}$, we conclude that $a = bc \in I$. Thus I is a weakly primary ideal of R .

(2) \Rightarrow (1) It is clear by Theorem 1(2). \square

Recall that a commutative ring R is called a *chained* ring if for every $x, y \in R$, we have $x \mid y$ or $y \mid x$. Every chained ring is divided. So, if R is a reduced chained ring, then a proper ideal I of R is a weakly 1-absorbing primary ideal if and only if it is a weakly primary ideal of R .

Theorem 8. *Let R be a Dedekind domain and I be a nonzero proper ideal of R . Then I is a weakly 1-absorbing primary ideal of R if and only if \sqrt{I} is a prime ideal of R .*

Proof. Suppose that I is a weakly 1-absorbing primary ideal of R . Then \sqrt{I} is a prime ideal of R by Theorem 3. The converse follows from [10, Theorem 14]. \square

Let R be a commutative ring with $1 \neq 0$. If an ideal of R contained in a finite union of ideals must be contained in one of those ideals, then R is said to be a *u-ring* [13]. In the next theorem, we give some characterizations of weakly 1-absorbing primary ideals in *u-rings*.

Theorem 9. *Let R be a commutative u-ring, and I a proper ideal of R . Then the following statements are equivalent.*

- (1) I is a weakly 1-absorbing primary ideal of R .
- (2) For every nonunit elements $a, b \in R$ with $ab \notin I$, $(I : ab) = (0 : ab)$ or $(I : ab) \subseteq \sqrt{I}$.
- (3) For every nonunit element $a \in R$ and every ideal I_1 of R with $I_1 \not\subseteq \sqrt{I}$, if $(I : aI_1)$ is a proper ideal of R , then $(I : aI_1) = (\{0\} : aI_1)$ or $(I : aI_1) \subseteq (I : a)$.
- (4) For all ideals I_1, I_2 of R with $I_1 \not\subseteq \sqrt{I}$, if $(I : I_1I_2)$ is a proper ideal of R , then $(I : I_1I_2) = (\{0\} : I_1I_2)$ or $(I : I_1I_2) \subseteq (I : I_2)$.
- (5) For all ideals I_1, I_2, I_3 of R with $0 \neq I_1I_2I_3 \subseteq I$, $I_1I_2 \subseteq I$ or $I_3 \subseteq \sqrt{I}$.

Proof. (1) \Rightarrow (2) Suppose that I is a weakly 1-absorbing primary ideal of R , $ab \notin I$ for some nonunit elements $a, b \in R$, and $c \in (I : ab)$. Then $abc \in I$. Since $ab \notin I$, c is nonunit. If $abc = 0$, then $c \in (0 : ab)$. Assume that $0 \neq abc \in I$. Since I is weakly 1-absorbing primary, we have $c \in \sqrt{I}$. Hence we conclude that $(I : ab) \subseteq (0 : ab) \cup \sqrt{I}$. Since R is a u-ring, we obtain that $(I : ab) = (0 : ab)$ or $(I : ab) \subseteq \sqrt{I}$.

(2) \Rightarrow (3) If $aI_1 \subseteq I$, then we are done. Suppose that $aI_1 \not\subseteq I$ for some nonunit element $a \in R$ and $c \in (I : aI_1)$. It is clear that c is nonunit. Then $acI_1 \subseteq I$. Now $I_1 \subseteq (I : ac)$. If $ac \in I$, then $c \in (I : a)$. Suppose that $ac \notin I$. Hence $(I : ac) = (0 : ac)$ or $(I : ac) \subseteq \sqrt{I}$ by (2). Thus $I_1 \subseteq (0 : ac)$ or $I_1 \subseteq \sqrt{I}$. Since $I_1 \not\subseteq \sqrt{I}$ by hypothesis, we conclude that $I_1 \subseteq (0 : ac)$; i.e. $c \in (\{0\} : aI_1)$. Thus $(I : aI_1) \subseteq (\{0\} : aI_1) \cup (I : a)$. Since R is a u-ring, we have $(I : aI_1) = (\{0\} : aI_1)$ or $(I : aI_1) \subseteq (I : a)$.

(3) \Rightarrow (4) If $I_1 \subseteq \sqrt{I}$, then we are done. Suppose that $I_1 \not\subseteq \sqrt{I}$ and $c \in (I : I_1I_2)$. Then $I_2 \subseteq (I : cI_1)$. Since $(I : I_1I_2)$ is proper, c is a nonunit. Hence $I_2 \subseteq (\{0\} : cI_1)$ or $I_2 \subseteq (I : c)$ by (3). If $I_2 \subseteq (\{0\} : cI_1)$, then $c \in (\{0\} : I_1I_2)$. If $I_2 \subseteq (I : c)$, then $c \in (I : I_2)$. So, $(I : I_1I_2) \subseteq (\{0\} : I_1I_2) \cup (I : I_2)$, which implies that $(I : I_1I_2) = (\{0\} : I_1I_2)$ or $(I : I_1I_2) \subseteq (I : I_2)$, as needed.

(4) \Rightarrow (5) It is clear.

(5) \Rightarrow (1) Let $a, b, c \in R$ be nonunit elements and $0 \neq abc \in I$. Put $I_1 = aR$, $I_2 = bR$, and $I_3 = cR$. Then (1) is now clear by (5). \square

Definition 2. *Let I be a weakly 1-absorbing primary ideal of a commutative ring R and a, b, c be nonunit elements of R . We call (a, b, c) a 1-triple-zero of I if $abc = 0$, $ab \notin I$, and $c \notin \sqrt{I}$.*

Observe that if I is a weakly 1-absorbing primary ideal of a commutative ring R that is not 1-absorbing primary, then there exists a 1-triple-zero (a, b, c) of I for some nonunit elements $a, b, c \in R$.

Theorem 10. *Let I be a weakly 1-absorbing primary ideal of a commutative ring R and (a, b, c) be a 1-triple-zero of I .*

- (1) $abI = \{0\}$.
- (2) If $a, b \notin (I : c)$, then $bcI = acI = aI^2 = bI^2 = cI^2 = \{0\}$.
- (3) If $a, b \notin (I : c)$, then $I^3 = \{0\}$.

Proof. (1) Suppose that $abI \neq \{0\}$. Then $abx \neq 0$ for some nonunit $x \in I$. Hence $0 \neq ab(c+x) \in I$. Since $ab \notin I$, $(c+x)$ is nonunit element of R . Since I is a weakly 1-absorbing primary ideal of R and $ab \notin I$, we conclude that $(c+x) \in \sqrt{I}$. Since $x \in I$, we have $c \in \sqrt{I}$, a contradiction. Thus $abI = \{0\}$.

(2) Suppose that $bcI \neq 0$. Then $bcy \neq 0$ for some nonunit element $y \in I$. Hence $0 \neq bcy = b(a+y)c \in I$. Since $b \notin (I : c)$, we conclude that $a+y$ is a nonunit element of R . Since I is a weakly 1-absorbing primary ideal of R , $ab \notin I$, and $by \in I$, we conclude that $b(a+y) \notin I$, and hence $c \in \sqrt{I}$, a contradiction. Thus $bcI = \{0\}$. We show that $acI = \{0\}$. Suppose that $acI \neq \{0\}$. Then $acy \neq 0$ for some nonunit element $y \in I$. Hence $0 \neq acy = a(b+y)c \in I$. Since $a \notin (I : c)$, we conclude that $b+y$ is a nonunit element of R . Since I is a weakly 1-absorbing primary ideal of R , $ab \notin I$, and $ay \in I$, we conclude that $a(b+y) \notin I$, and hence $c \in \sqrt{I}$, a contradiction. Thus $acI = \{0\}$. Now we prove that $aI^2 = \{0\}$. Suppose that $axy \neq 0$ for some $x, y \in I$. Since $abI = \{0\}$ by (1) and $acI = \{0\}$ by (2), $0 \neq axy = a(b+x)(c+y) \in I$. Since $ab \notin I$, we conclude that $c+y$ is a nonunit element of R . Since $a \notin (I : c)$, we conclude that $b+x$ is a nonunit element of R . Since I is a weakly 1-absorbing primary ideal of R , we have $a(b+x) \in I$ or $(c+y) \in \sqrt{I}$. Since $x, y \in I$, we conclude that $ab \in I$ or $c \in \sqrt{I}$, a contradiction. Thus $aI^2 = \{0\}$. We show $bI^2 = \{0\}$. Suppose that $bxy \neq 0$ for some $x, y \in I$. Since $abI = \{0\}$ by (1) and $bcI = \{0\}$ by (2), $0 \neq bxy = b(a+x)(c+y) \in I$. Since $ab \notin I$, we conclude that $c+y$ is a nonunit element of R . Since $b \notin (I : c)$, we conclude that $a+x$ is a nonunit element of R . Since I is a weakly 1-absorbing primary ideal of R , we have $b(a+x) \in I$ or $(c+y) \in \sqrt{I}$. Since $x, y \in I$, we conclude that $ab \in I$ or $c \in \sqrt{I}$, a contradiction. Thus $bI^2 = \{0\}$. We show $cI^2 = \{0\}$. Suppose that $cxy \neq 0$ for some $x, y \in I$. Since $acI = bcI = \{0\}$ by (2), $0 \neq cxy = (a+x)(b+y)c \in I$. Since $a, b \notin (I : c)$, we conclude that $a+x$ and $b+y$ are nonunit elements of R . Since I is a weakly 1-absorbing primary ideal of R , we have $(a+x)(b+y) \in I$ or $c \in \sqrt{I}$. Since $x, y \in I$, we conclude that $ab \in I$ or $c \in \sqrt{I}$, a contradiction. Thus $cI^2 = \{0\}$.

(3) Assume that $xyz \neq 0$ for some $x, y, z \in I$. Then $0 \neq xyz = (a+x)(b+y)(c+z) \in I$ by (1) and (2). Since $ab \notin I$, we conclude $c+z$ is a nonunit element of R . Since $a, b \notin (I : c)$, we conclude that $a+x$ and $b+y$ are nonunit elements of R . Since I is a weakly 1-absorbing primary ideal of R , we have $(a+x)(b+y) \in I$ or $c+z \in \sqrt{I}$. Since $x, y, z \in I$, we conclude that $ab \in I$ or $c \in \sqrt{I}$, a contradiction. Thus $I^3 = \{0\}$. \square

Theorem 11. (1) *Let I be a weakly 1-absorbing primary ideal of a commutative reduced ring R . Suppose that I is not a 1-absorbing primary ideal of R and (a, b, c) is a 1-triple-zero of I such that $a, b \notin (I : c)$. Then $I = \{0\}$.*

- (2) Let I be a nonzero weakly 1-absorbing primary ideal of a reduced ring R . Suppose that I is not a 1-absorbing primary ideal of R and (a, b, c) is a 1-triple-zero of I . Then $ac \in I$ or $bc \in I$.

Proof. (1) Since $a, b \in (I : c)$, then $I^3 = \{0\}$ by Theorem 10(3). Since R is reduced, we conclude that $I = \{0\}$.

(2) Suppose that neither $ac \in I$ nor $bc \in I$. Then $I = \{0\}$ by (1), a contradiction since I is a nonzero ideal of R by hypothesis. Hence if (a, b, c) is a 1-triple-zero of I , then $ac \in I$ or $bc \in I$. \square

Theorem 12. Let I be a weakly 1-absorbing primary ideal of a commutative ring R . If I is not a weakly primary ideal of R , then there exist an irreducible element $x \in R$ and a nonunit element $y \in R$ such that $xy \in I$, but neither $x \in I$ nor $y \in \sqrt{I}$. Furthermore, if $ab \in I$ for some nonunit elements $a, b \in R$ such that neither $a \in I$ nor $b \in \sqrt{I}$, then a is an irreducible element of R .

Proof. Suppose that I is not a weakly primary ideal of R . Then there exist nonunit elements $x, y \in R$ such that $0 \neq xy \in I$ with $x \notin I$, $y \notin \sqrt{I}$. Suppose that x is not an irreducible element of R . Then $x = cd$ for some nonunit elements $c, d \in R$. Since $0 \neq xy = cdy \in I$, I is weakly 1-absorbing primary, and $y \notin \sqrt{I}$, we conclude that $x = cd \in I$, a contradiction. Hence x is an irreducible element of R . \square

In general, the intersection of a family of weakly 1-absorbing primary ideals need not be a weakly 1-absorbing primary ideal. Indeed, consider the ring $R = \mathbb{Z}_{12}$. Then $I = (2)$ and $J = (3)$ are clearly weakly 1-absorbing primary ideals of R , but $I \cap J = \{0, 6\}$ is not a weakly 1-absorbing primary ideal of R (since $0 \neq 3 \cdot 3 \cdot 2 \in I \cap J$, but neither $3 \cdot 3 \in I \cap J$ nor $2 \in \sqrt{I \cap J}$). However, we have the following result.

Proposition 1. Let $\{I_i \mid i \in \Lambda\}$ be a finite collection of weakly 1-absorbing primary ideals of a commutative ring R such that $Q = \sqrt{I_i} = \sqrt{I_j}$ for every distinct $i, j \in \Lambda$. Then $I = \bigcap_{i \in \Lambda} I_i$ is a weakly 1-absorbing primary ideal of R .

Proof. Suppose that $0 \neq abc \in I = \bigcap_{i \in \Lambda} I_i$ for nonunit elements a, b, c of R and $ab \notin I$. Then for some $k \in \Lambda$, $0 \neq abc \in I_k$ and $ab \notin I_k$. This implies that $c \in \sqrt{I_k} = Q = \sqrt{I}$. \square

Proposition 2. Let I be a weakly 1-absorbing primary ideal of a commutative ring R and c be a nonunit element of $R \setminus I$. Then $(I : c)$ is a weakly primary ideal of R .

Proof. Suppose that $0 \neq ab \in (I : c)$ for some nonunit $c \in R \setminus I$ and assume that $a \notin (I : c)$. Hence b is a nonunit element of R . If a is a unit of R , then $b \in (I : c) \subseteq \sqrt{(I : c)}$ and we are done. So assume that a is a nonunit element of R . Since $0 \neq abc = acb \in I$, $ac \notin I$, and I is a weakly 1-absorbing primary ideal of R , we conclude that $b \in \sqrt{I} \subseteq \sqrt{(I : c)}$. Thus $(I : c)$ is a weakly primary ideal of R . \square

The next theorem gives a characterization for weakly 1-absorbing primary ideals of $R = R_1 \times R_2$, where R_1 and R_2 are commutative rings with identity that are not fields.

Theorem 13. Let R_1 and R_2 be commutative rings with identity that are not fields, $R = R_1 \times R_2$, and I be a nonzero proper ideal of R . Then the following statements are equivalent.

- (1) I is a weakly 1-absorbing primary ideal of R .
- (2) $I = I_1 \times R_2$ for some primary ideal I_1 of R_1 or $I = R_1 \times I_2$ for some primary ideal I_2 of R_2 .
- (3) I is a 1-absorbing primary ideal of R .
- (4) I is a primary ideal of R .

Proof. (1) \Rightarrow (2) Suppose that I is a weakly 1-absorbing primary ideal of R . Then I is of the form $I_1 \times I_2$ for some ideals I_1 and I_2 of R_1 and R_2 , respectively. Assume that both I_1 and I_2 are proper. Since I is a nonzero ideal of R , we conclude that $I_1 \neq \{0\}$ or $I_2 \neq \{0\}$. We may assume that $I_1 \neq \{0\}$. Let $0 \neq c \in I_1$. Then $0 \neq (1, 0)(1, 0)(c, 1) = (c, 0) \in I_1 \times I_2$. This implies that $(1, 0)(1, 0) \in I_1 \times I_2$ or $(c, 1) \in \sqrt{I_1 \times I_2} = \sqrt{I_1} \times \sqrt{I_2}$, that is $I_1 = R_1$ or $I_2 = R_2$, a contradiction. Thus either I_1 or I_2 is a proper ideal. Without loss of generality, assume that $I = I_1 \times R_2$ for some proper ideal I_1 of R_1 . We show that I_1 is a primary ideal of R_1 . Let $ab \in I_1$ for some $a, b \in R_1$. We can assume that a and b are nonunit elements of R_1 . Since R_2 is not a field, there exists a nonunit nonzero element $x \in R_2$. Then $0 \neq (a, 1)(1, x)(b, 1) \in I_1 \times R_2$ which implies that either $(a, 1)(1, x) \in I_1 \times R_2$ or $(b, 1) \in \sqrt{I_1 \times R_2} = \sqrt{I_1} \times R_2$; i.e., $a \in I_1$ or $b \in \sqrt{I_1}$.

(2) \Rightarrow (3) Since I is a primary ideal of R , I is a 1-absorbing primary ideal of R by [10, Theorem 1(1)].

(3) \Rightarrow (4) Since I a 1-absorbing primary ideal of R and R is not a quasilocal ring, we conclude that I is a primary ideal of R by [10, Theorem 3].

(4) \Rightarrow (1) It is clear. \square

Theorem 14. *Let R_1, \dots, R_n be commutative rings with $1 \neq 0$ for some $2 \leq n < \infty$, and let $R = R_1 \times \dots \times R_n$. Then the following statements are equivalent.*

- (1) Every proper ideal of R is a weakly 1-absorbing primary ideal of R .
- (2) $n = 2$ and R_1, R_2 are fields.

Proof. (1) \Rightarrow (2) Suppose that every proper ideal of R is a weakly 1-absorbing primary ideal. Without loss of generality, we may assume that $n = 3$. Then $I = R_1 \times \{0\} \times \{0\}$ is a weakly 1-absorbing primary ideal of R . However, for a nonzero $a \in R_1$, we have $(0, 0, 0) \neq (1, 0, 1)(1, 0, 1)(a, 1, 0) = (a, 0, 0) \in I$, but neither $(1, 0, 1)(1, 0, 1) \in I$ nor $(a, 1, 0) \in \sqrt{I}$, a contradiction. Thus $n = 2$. Assume that R_1 is not a field. Then there exists a nonzero proper ideal A of R_1 . Hence $I = A \times \{0\}$ is a weakly 1-absorbing primary ideal of R . However, for a nonzero $a \in A$, we have $(0, 0) \neq (1, 0)(1, 0)(a, 1) = (a, 0) \in I$, but neither $(1, 0)(1, 0) \in I$ nor $(a, 1) \in \sqrt{I}$, a contradiction. Similarly, one can easily show that R_2 is a field. Hence $n = 2$ and R_1, R_2 are fields.

(2) \Rightarrow (1) Suppose that $n = 2$ and R_1, R_2 are fields. Then R has exactly three proper ideals, i.e., $\{(0, 0)\}$, $\{0\} \times R_2$, and $R_1 \times \{0\}$ are the only proper ideals of R . Hence it is clear that each proper ideal of R is a weakly 1-absorbing primary ideal of R . \square

Since every ring that is a product of a finite number of fields is a von Neumann regular ring, in light of Theorem 4 and Theorem 14, we have the following result.

Corollary 2. *Let R_1, \dots, R_n be commutative rings with $1 \neq 0$ for some $2 \leq n < \infty$, and let $R = R_1 \times \dots \times R_n$. Then the following statements are equivalent.*

- (1) Every proper ideal of R is a weakly 1-absorbing primary ideal of R .
- (2) Every proper ideal of R is a weakly primary ideal of R .

- (3) $n = 2$ and R_1, R_2 are fields, and hence $R = R_1 \times R_2$ is a von Neumann regular ring.

Theorem 15. *Let R_1 and R_2 be commutative rings and $f : R_1 \rightarrow R_2$ be a ring homomorphism with $f(1) = 1$.*

- (1) Suppose that f is injective, $f(a)$ is a nonunit element of R_2 for every nonunit element $a \in R_1$, and J is a weakly 1-absorbing primary ideal of R_2 . Then $f^{-1}(J)$ is a weakly 1-absorbing primary ideal of R_1 .
- (2) If f is an epimorphism and I is a weakly 1-absorbing primary ideal of R_1 such that $\text{Ker}(f) \subseteq I$, then $f(I)$ is a weakly 1-absorbing primary ideal of R_2 .

Proof. (1) Since $f(1) = 1$, $f^{-1}(J)$ is a proper ideal of R_1 . Let $0 \neq abc \in f^{-1}(J)$ for some nonunit elements $a, b, c \in R$. Since $\text{Ker}(f) = 0$, we have $0 \neq f(abc) = f(a)f(b)f(c) \in J$, where $f(a), f(b), f(c)$ are nonunit elements of R_2 by hypothesis. Hence $f(a)f(b) \in J$ or $f(c) \in \sqrt{J}$. Hence $ab \in f^{-1}(J)$ or $c \in \sqrt{f^{-1}(J)} = f^{-1}(\sqrt{J})$. Thus $f^{-1}(J)$ is a weakly 1-absorbing primary ideal of R_1 .

(2) Let $0 \neq xyz \in f(I)$ for some nonunit elements $x, y, z \in R$. Since f is onto, there exist nonunit elements $a, b, c \in I$ such that $x = f(a), y = f(b), z = f(c)$. Then $f(abc) = f(a)f(b)f(c) = xyz \in f(I)$. Since $\text{Ker}(f) \subseteq I$, we have $0 \neq abc \in I$. Hence $ab \in I$ or $c \in \sqrt{I}$. Thus $xy \in f(I)$ or $z \in f(\sqrt{I})$. Since f is onto and $\text{Ker}(f) \subseteq I$, we have $f(\sqrt{I}) = \sqrt{f(I)}$. Thus we are done. \square

The following example shows that the hypothesis in Theorem 15(1) is crucial.

Example 2. ([10, Example 1]) *Let $A = K[x, y]$, where K is a field, $M = (x, y)A$, and $B = A_M$. Note that B is a quasilocal ring with maximal ideal M_M . Then $I = xM_M = (x^2, xy)B$ is a 1-absorbing primary ideal of B (see [10, Theorem 5]) and $\sqrt{I} = xB$. However $xy \in I$, but neither $x \in I$ nor $y \in \sqrt{I}$. Thus I is not a primary ideal of B . Let $f : B \times B \rightarrow B$ such that $f(x, y) = x$. Then f is a ring homomorphism from $B \times B$ onto B such that $f(1, 1) = 1$. However, $(1, 0)$ is a nonunit element of $B \times B$ and $f(1, 0) = 1$ is a unit of B . Thus f does not satisfy the hypothesis of Theorem 15(1). Now $f^{-1}(I) = I \times B$ is not a weakly 1-absorbing primary ideal of $B \times B$ by Theorem 13.*

Theorem 16. *Let I be a proper ideal of a commutative ring R .*

- (1) *If J is a proper ideal of R with $J \subseteq I$ and I is a weakly 1-absorbing primary ideal of R , then I/J is a weakly 1-absorbing primary ideal of R/J .*
- (2) *Let J be a proper ideal of R with $J \subseteq I$ such that $a + J$ is a nonunit element of R/J for every nonunit $a \in R$. If J is a 1-absorbing primary ideal of R and I/J is a weakly 1-absorbing primary ideal of R/J , then I is a 1-absorbing primary ideal of R .*
- (3) *If $\{0\}$ is a 1-absorbing primary ideal of R and I is a weakly 1-absorbing primary ideal of R , then I is a 1-absorbing primary ideal of R .*
- (4) *Let J be a proper ideal of R with $J \subseteq I$ such that $a + J$ is a nonunit element of R/J for every nonunit $a \in R$. If J is a weakly 1-absorbing primary ideal of R and I/J is a weakly 1-absorbing primary ideal of R/J , then I is a weakly 1-absorbing primary ideal of R .*

Proof. (1) Consider the natural epimorphism $\pi : R \rightarrow R/J$. Then $\pi(I) = I/J$. So we are done by Theorem 15 (2).

(2) Suppose that $abc \in I$ for some nonunit elements $a, b, c \in R$. If $abc \in J$, then $ab \in J \subseteq I$ or $c \in \sqrt{J} \subseteq \sqrt{I}$ as J is a 1-absorbing primary ideal of R . Now assume that $abc \notin J$. Then $J \neq (a+J)(b+J)(c+J) \in I/J$, where $a+J, b+J, c+J$ are nonunit elements of R/J by hypothesis. Thus $(a+J)(b+J) \in I/J$ or $(c+J) \in \sqrt{I/J}$. Hence $ab \in I$ or $c \in \sqrt{I}$.

(3) The proof follows from (2).

(4) Suppose that $0 \neq abc \in I$ for some nonunit elements $a, b, c \in R$. If $abc \in J$, then $ab \in J \subseteq I$ or $c \in \sqrt{J} \subseteq \sqrt{I}$ as J is a weakly 1-absorbing primary ideal of R . Now assume that $abc \notin J$. Then $J \neq (a+J)(b+J)(c+J) \in I/J$, where $a+J, b+J, c+J$ are nonunit elements of R/J by hypothesis. Thus $(a+J)(b+J) \in I/J$ or $(c+J) \in \sqrt{I/J}$. Hence $ab \in I$ or $c \in \sqrt{I}$. \square

In the following remark, we give the correct version of [10, Theorem 17(1), Corollary 3, and Corollary 4].

Remark 1. *Mohammed Tamekkante pointed out to the first-named author that in [10], we overlooked the fact that if $f : R_1 \rightarrow R_2$ is a ring homomorphism such that $f(1) = 1$, then it is possible that $f(a) \in U(R_2)$ for some nonunit element $a \in R_1$. Overlooking this fact caused a problem in the proof of [10, Theorem 17(1), Corollary 3, and Corollary 4]. We state the correct version of [10, Theorem 17(1), Corollary 3, and Corollary 4].*

- (1) ([10, Theorem 17(1)]). *Let R_1 and R_2 be commutative rings and $f : R_1 \rightarrow R_2$ be a ring homomorphism with $f(1) = 1$ such that if R_2 is a quasilocal ring, then $f(a)$ is a nonunit element of R_2 for every nonunit element $a \in R_1$. If J is a 1-absorbing primary ideal of R_2 , then $f^{-1}(J)$ is a 1-absorbing primary ideal of R_1 . (Note that if R_2 is not a quasilocal ring, then J is primary by [10, Theorem 3], and hence $f^{-1}(J)$ is a primary ideal of R_1 . Since every primary ideal of a commutative ring A is a 1-absorbing primary ideal of A , we conclude that $f^{-1}(J)$ is a 1-absorbing primary ideal of R_1 .)*
- (2) ([10, Corollary 3]). *Let I and J be proper ideals of a commutative ring R with $I \subseteq J$. If J is a 1-absorbing primary ideal of R , then J/I is a 1-absorbing primary ideal of R/I . Furthermore, assume that if R/I is a quasilocal ring, then $a+I$ is a nonunit element of R/I for every nonunit $a \in R$. If J/I is a 1-absorbing primary ideal of R/I , then J is a 1-absorbing primary ideal of R .*
- (3) ([10, Corollary 4]). *Let R be a commutative ring and $A = R[x]$. Then a proper ideal I of R is a 1-absorbing primary ideal of R if and only if $(I[x] + xA)/xA$ is a 1-absorbing primary ideal of A/xA . (The claim is clear since R is ring-isomorphic to A/xA .)*

Note that Example 2 shows that the hypothesis in (1) is crucial.

Theorem 17. *Let S be a multiplicatively closed subset of a commutative ring R , and I be proper ideal of R .*

- (1) *If I is a weakly 1-absorbing primary ideal of R such that $I \cap S = \emptyset$, then $S^{-1}I$ is a weakly 1-absorbing primary ideal of $S^{-1}R$.*
- (2) *If $S^{-1}I$ is a weakly 1-absorbing primary ideal of $S^{-1}R$ such that $S \cap Z(R) = \emptyset$ and $S \cap Z_I(R) = \emptyset$, then I is a weakly 1-absorbing primary ideal of R .*

Proof. (1) Suppose that $0 \neq \frac{a}{s_1} \frac{b}{s_2} \frac{c}{s_3} \in S^{-1}I$ for some nonunit $a, b, c \in R \setminus S$, $s_1, s_2, s_3 \in S$ and $\frac{a}{s_1} \frac{b}{s_2} \notin S^{-1}I$. Then $0 \neq uabc \in I$ for some $u \in S$. Since I is

weakly 1-absorbing primary and $uab \notin I$, we conclude $c \in \sqrt{I}$. Thus $\frac{c}{s_3} \in S^{-1}\sqrt{I} = \sqrt{S^{-1}I}$. Thus $S^{-1}I$ is a weakly 1-absorbing primary ideal of $S^{-1}R$.

(2) Suppose that $0 \neq abc \in I$ for some nonunit elements $a, b, c \in R$. Hence $0 \neq \frac{abc}{1} = \frac{a}{1}\frac{b}{1}\frac{c}{1} \in S^{-1}I$ as $S \cap Z(R) = \emptyset$. Since $S^{-1}I$ is weakly 1-absorbing primary, we have either $\frac{a}{1} \in S^{-1}I$ or $\frac{c}{1} \in \sqrt{S^{-1}I} = S^{-1}\sqrt{I}$. If $\frac{a}{1} \in S^{-1}I$, then $uab \in I$ for some $u \in S$. Since $S \cap Z_I(R) = \emptyset$, we conclude that $ab \in I$. If $\frac{c}{1} \in S^{-1}\sqrt{I}$, then $(tc)^n \in I$ for some positive integer $n \geq 1$ and $t \in S$. Since $t^n \notin Z_I(R)$, we have $c^n \in I$, i.e., $c \in \sqrt{I}$. Thus I is a weakly 1-absorbing primary ideal of R . \square

Definition 3. Let I be a weakly 1-absorbing primary ideal of a commutative ring R and $I_1I_2I_3 \subseteq I$ for some proper ideals I_1, I_2, I_3 of R . If (a, b, c) is not a 1-triple-zero of I for every $a \in I_1, b \in I_2, c \in I_3$, then we call I a free 1-triple-zero with respect to $I_1I_2I_3$.

Theorem 18. Let I be a weakly 1-absorbing primary ideal of a commutative ring R and J be a proper ideal of R with $abJ \subseteq I$ for some $a, b \in R$. If (a, b, j) is not a 1-triple-zero of I for all $j \in J$ and $ab \notin I$, then $J \subseteq \sqrt{I}$.

Proof. Suppose that $J \not\subseteq \sqrt{I}$. Then there exists $c \in J \setminus \sqrt{I}$. Then $abc \in abJ \subseteq I$. If $abc \neq 0$, then it contradicts our assumption that $ab \notin I$ and $c \notin \sqrt{I}$. Thus $abc = 0$. Since (a, b, c) is not a 1-triple-zero of I and $ab \notin I$, we conclude that $c \in \sqrt{I}$, a contradiction. Thus $J \subseteq \sqrt{I}$. \square

Theorem 19. Let I be a weakly 1-absorbing primary ideal of a commutative ring R and $\{0\} \neq I_1I_2I_3 \subseteq I$ for some proper ideals I_1, I_2, I_3 of R . If I is free 1-triple-zero with respect to $I_1I_2I_3$, then $I_1I_2 \subseteq I$ or $I_3 \subseteq \sqrt{I}$.

Proof. Suppose that I is free 1-triple-zero with respect to $I_1I_2I_3$, and $\{0\} \neq I_1I_2I_3 \subseteq I$. Assume that $I_1I_2 \not\subseteq I$. Then there exist $a \in I_1, b \in I_2$ such that $ab \notin I$. Since I is a free 1-triple-zero with respect to $I_1I_2I_3$, we conclude that (a, b, c) is not a 1-triple-zero of I for all $c \in I_3$. Thus $I_3 \subseteq \sqrt{I}$ by Theorem 18. \square

Acknowledgement. We would like to thank Mohammed Tamekkante for pointing out a problem in [10, Theorem 17(1)]. Also, we are so grateful to the referee for his/her great effort in proofreading the manuscript.

REFERENCES

- [1] D. D. Anderson and M. Batanieh, Generalizations of prime ideals, *Comm. Algebra*, **36** (2008), 686–696.
- [2] D.D. Anderson and E. Smith, Weakly prime ideals, *Houston J. Math.*, **29** (4) (2003), 831–840.
- [3] D.F. Anderson and A. Badawi, On n -absorbing ideals of commutative rings, *Comm. Algebra*, **39** (2011), 1646–1672.
- [4] D.F. Anderson and A. Badawi, Von Neumann regular and related elements in commutative rings, *Algebra Colloq.*, **19**(SPL. ISS. 1) (2012), 1017–1040.
- [5] S. E. Atani and F. Farzalipour, On weakly primary ideals, *Georgian Math. J.*, **12**(3) (2005), 423–429.
- [6] A.Badawi, On 2-absorbing ideals of commutative rings, *Bull. Austral. Math. Soc.*, **75** (2007), 417–429.
- [7] A. Badawi and A. Y. Darani, On weakly 2-absorbing ideals of commutative rings, *Houston J. Math.* **39** (2013), no. 2, 441–452.
- [8] A. Badawi, Ü. Tekir and E. Yetkin, On 2-absorbing primary ideals in commutative rings, *Bull. Korean Math. Soc.*, **51** (4) (2014), 1163–1173.

- [9] A. Badawi, Ü. Tekir and E. Yetkin, On weakly 2-absorbing primary ideals in commutative rings, J. Korean Math. Soc., **52**(1) (2015), 97-111.
- [10] A. Badawi and E. Yetkin Çelikel, On 1-absorbing primary ideals of a commutative ring, J. Algebra Appl., **19**(6) (2020), (12 pages).
- [11] A. Badawi, D. Sonmez and G. Yesilot, On Weakly δ -Semiprimary Ideals of Commutative Rings, Algebra Colloq., **25**(3) (2018), 387-398.
- [12] R. W. Gilmer, Rings in which semiprimary ideals are primary, Pacific J. Math., **12**(4) (1962) 1273-1276.
- [13] P. Quartararo and H. S. Butts, Finite unions of ideals and modules, Proc. Amer. Math. Soc., **52** (1975), 91-96.
- [14] O. Zariski and Pierre Samuel, Commutative Algebra. V.I. (Princeton, 1958).

DEPARTMENT OF MATHEMATICS & STATISTICS, THE AMERICAN UNIVERSITY OF SHARJAH, P.O. BOX 26666, SHARJAH, UNITED ARAB EMIRATES.

Email address: abadawi@aus.edu

DEPARTMENT OF ELECTRICAL-ELECTRONICS ENGINEERING, FACULTY OF ENGINEERING, HASAN KALYONCU UNIVERSITY, GAZIANTEP, TURKEY.

Email address: yetkinece@gmail.com; ece.celikel@hku.edu.tr