

ON 1-ABSORBING PRIMARY IDEAL OF A COMMUTATIVE RING (CORRECTION TO THEOREM 18 IS ADDED)

AYMAN BADAWI AND ECE YETKIN CELIKEL

ABSTRACT. Let R be a commutative ring with nonzero identity. In this paper, we introduce the concept of 1-absorbing primary ideals in commutative rings. A proper ideal I of R is called a 1-absorbing primary ideal of R if whenever nonunit elements $a, b, c \in R$ and $abc \in I$, then $ab \in I$ or $c \in \sqrt{I}$. Some properties of 1-absorbing primary ideals are investigated. For example, we show that if R admits a 1-absorbing primary ideal that is not a primary ideal, then R is a quasilocal ring. We give an example of a 1-absorbing primary ideal of R that is not a primary ideal of R . We show that if a ring R is not a quasilocal, then a proper ideal I of R is a 1-absorbing primary ideal of R if and only if I is a primary ideal. We show that if R is a Noetherian domain, then R is a Dedekind domain if and only if every nonzero proper 1-absorbing primary ideal of R is of the form P^n for some nonzero prime ideal P of R and a positive integer $n \geq 1$. We show that a proper ideal I of R is a 1-absorbing primary ideal of R if and only if whenever $I_1 I_2 I_3 \subseteq I$ for some proper ideals I_1, I_2, I_3 of R , then $I_1 I_2 \subseteq I$ or $I_3 \subseteq \sqrt{I}$.

1. INTRODUCTION

Throughout this paper all rings are commutative with $1 \neq 0$. Let R be a commutative ring. If R has exactly one maximal ideal, then R is called a *quasilocal* ring. An ideal I of R is said to be proper if $I \neq R$. Let I be a proper ideal of a commutative ring R . Then the radical of I is denoted by $\sqrt{I} = \{r \in R \mid r^n \in I \text{ for some positive integer } n \geq 1\}$ and the set of zero divisor elements with respect to I is denoted by $Z_I(R) = \{r \in R \mid rs \in I \text{ for some } s \in R \setminus I\}$.

Since prime ideals have an important role in the theory of commutative rings, there are several ways to generalize the concept of prime ideals. Badawi generalized the concept of prime ideals in [3]. We recall from [3] that a nonzero proper ideal I of R is said to be a 2-absorbing ideal of R if whenever $a, b, c \in R$ and $abc \in I$, then either $ab \in I$ or $ac \in I$ or $bc \in I$. Anderson and Badawi [2] generalized the notion of 2-absorbing ideals to n -absorbing ideals. A proper ideal I of R is called n -absorbing ideal if whenever $x_1 \cdots x_{n+1} \in I$ for $x_1, \dots, x_{n+1} \in R$, then there are n of the x_i 's whose product is in I . Recall from [4] that a proper ideal I of R is called a 2-absorbing primary ideal of R if whenever $a, b, c \in R$ with $abc \in I$, then $ab \in I$ or $ac \in \sqrt{I}$ or $bc \in \sqrt{I}$.

In this paper, we introduce the concept of 1-absorbing primary ideals of commutative rings. A proper ideal I of a commutative ring R is called a 1-absorbing primary ideal of R if whenever nonunit elements $a, b, c \in R$ and $abc \in I$, then

2000 *Mathematics Subject Classification.* Primary 13A15, 13F05; Secondary 05A15, 13G05.

Key words and phrases. prime ideal, primary ideal, 1-absorbing primary ideal, 2-absorbing primary ideal, 2-absorbing ideal, weakly prime, weakly primary ideal, weakly 2-absorbing primary ideal, weakly semiprime, n -absorbing ideal.

$ab \in I$ or $c \in \sqrt{I}$. We show that the following implications hold and none of them is revisable:

primary ideal \implies 1-absorbing primary ideal \implies 2-absorbing primary ideal.

Among many results in this paper. We give an example (Example 1) of a 1-absorbing primary ideal of R that is not a primary ideal of R , and another example (Example 2) of a 2-absorbing primary ideal of R that is not a 1-absorbing primary ideal of R . We show (Theorem 2) that if I is a 1-absorbing primary ideal of R , then \sqrt{I} is a prime ideal of R . We show (Theorem 3) if a ring R admits a 1-absorbing primary ideal of R that is not a primary ideal, then R is a quasilocal ring. We give a method (Theorem 6) to construct 1-absorbing primary ideals of commutative rings that are not primary ideals. We show (Theorem 7) that if a ring R is not a quasilocal, then a proper ideal I of R is a 1-absorbing primary ideal of R if and only if I is a primary ideal. We show (Theorem 16) that if R is a Noetherian domain, then R is a Dedekind domain if and only if every nonzero proper 1-absorbing primary ideal of R is of the form P^n for some nonzero prime ideal P of R and a positive integer $n \geq 1$. We show (Theorem 21) that a proper ideal I of R is a 1-absorbing primary ideal of R if and only if whenever $I_1 I_2 I_3 \subseteq I$ for some proper ideals I_1, I_2, I_3 of R , then $I_1 I_2 \subseteq I$ or $I_3 \subseteq \sqrt{I}$.

For any undefined terminology see [5], [6], [7], and [8].

2. PROPERTIES OF 1-ABSORBING PRIMARY IDEALS

We remind the reader with the following definitions.

Definition 1. *Let I be a proper ideal of a commutative ring R .*

- (1) *We call I a 1-absorbing primary ideal of R if whenever nonunit elements $a, b, c \in R$ and $abc \in I$, then $ab \in I$ or $c \in \sqrt{I}$.*
- (2) ([4]) *We call I a 2-absorbing primary ideal of R if whenever $a, b, c \in R$ with $abc \in I$, then $ab \in I$ or $ac \in \sqrt{I}$ or $bc \in \sqrt{I}$.*

We start with the following trivial result, and hence we omit its proof.

Theorem 1. *Let I be a proper ideal of R . Then*

- (1) *If I is a primary ideal of R , then I is a 1-absorbing primary ideal of R .*
- (2) *If I is a 1-absorbing primary ideal of R , then I is a 2-absorbing primary ideal of R .*

The following is an example of a 1-absorbing primary ideal that is not a primary ideal.

Example 1. *Let $A = K[x, y]$, where K is a field, $M = (x, y)A$, and $R = A_M$. Note that R is a quasilocal ring with maximal ideal M_M . Then $I = xM_M = (x^2, xy)R$ is a 1-absorbing primary ideal of R (see Theorem 6) and $\sqrt{I} = xR$. However $xy \in I$, but neither $x \in I$ nor $y \in \sqrt{I}$. Thus I is not a primary ideal of R .*

The following is an example of a 2-absorbing primary ideal that is not a 1-absorbing primary ideal.

Example 2. *Let $R = \mathbb{Z}$. Consider the ideal $I = 12R$. Then I is a 2-absorbing primary ideal of R by Corollary 2.12 in [4]. However $2 \cdot 2 \cdot 3 \in I$, but neither $2 \cdot 2 \in I$ nor $3 \in \sqrt{I}$. Thus I is not a 1-absorbing primary ideal of R .*

Theorem 2. *Let I be a 1-absorbing primary ideal of a ring R . Then \sqrt{I} is a prime ideal of R .*

Proof. Let $xy \in \sqrt{I}$ for some $x, y \in R$. We may assume that x, y are nonunit elements of R . Let $n \geq 2$ be an even positive integer such that $(xy)^n \in I$. Then $n = 2m$ for some positive integer $m \geq 1$. Since $(xy)^n = x^n y^n = x^m x^m y^n \in I$ and I is a 1-absorbing primary ideal of R , we conclude that $x^m x^m = x^n \in I$ or $y^n \in I$. Hence $x \in \sqrt{I}$ or $y \in \sqrt{I}$. Thus \sqrt{I} is a prime ideal of R . \square

The following lemma is needed in the proof of our next result.

Lemma 1. *Let R be a ring. Suppose that for every nonunit element w of R and for every unit element u of R , we have $w + u$ is a unit element of R . Then R is a quasilocal ring.*

Proof. Suppose that R has at least two maximal ideals, say M_1, M_2 . Then $m_1 + m_2 = 1$ for some $m_1 \in M_1$ and $m_2 \in M_2$. Thus $1 - m_1 = m_2$ is a unit element of R , which is impossible since $m_2 \in M_2$. Thus R is a quasilocal ring. \square

In the following result, we show that if a ring R admits a 1-absorbing primary ideal that is not a primary ideal, then R is a quasilocal ring.

Theorem 3. *Suppose that a ring R admits a 1-absorbing primary ideal that is not a primary ideal. Then R is a quasilocal ring.*

Proof. Suppose that I is a 1-absorbing primary ideal of R that is not a primary ideal of R . Hence there exist nonunit elements $x, y \in R$ such that neither $x \in I$ nor $y \in \sqrt{I}$. Let w be a nonunit element of R . Since $wxy \in I$ and I is a 1-absorbing primary ideal of R and $y \notin \sqrt{I}$, we conclude that $wx \in I$. Let u be a unit element of R . Suppose that $w + u$ is a nonunit element of R . Since $(w + u)xy \in I$ and I is a 1-absorbing primary ideal of R and $y \notin \sqrt{I}$, we conclude that $(w + u)x = wx + ux \in I$. Since $wx \in I$, we conclude that $x \in I$, which is a contradiction. Thus $w + u$ is a unit element of R . Since for every nonunit element w of R and for every unit element u of R , we have $w + u$ is a unit element of R , we conclude that R is a quasilocal ring by Lemma 1. \square

Theorem 4. *Suppose that a ring R is not a quasilocal ring. Then a proper ideal I of R is a 1-absorbing primary ideal of R if and only if I is a primary ideal of R . In particular, if $R = R_1 \times R_2$ for some rings R_1 and R_2 , then a proper ideal I of R is a 1-absorbing primary ideal of R if and only if I is a primary ideal of R .*

Proof. If I is a primary ideal of R , then it is clear that I is a 1-absorbing primary ideal of R . Hence assume that I is a 1-absorbing primary ideal of R . Let $xy \in I$ for some $x, y \in R$. We may assume that x, y are nonunit elements of R . Suppose that $y \notin \sqrt{I}$. Since R is not a quasilocal ring, there exist a nonunit element $w \in R$ and a unit element $u \in R$ such that $w + u$ is a nonunit element of R by Lemma 1. Since $wxy \in I$ and I is a 1-absorbing primary ideal of R and $y \notin \sqrt{I}$, we conclude that $wx \in I$. Also, since $(w + u)xy \in I$ and I is a 1-absorbing primary ideal of R and $y \notin \sqrt{I}$, we conclude that $(w + u)x = wx + ux \in I$. Since $wx \in I$, we conclude that $x \in I$. Thus I is a primary ideal of R . \square

Let $R = R_1 \times R_2$, where R_1 and R_2 are commutative rings with $1 \neq 0$, and let J be a proper ideal of R . Then it is well-known that J is a primary ideal of R if and

only if $J = I \times R_2$ for some primary ideal I of R_1 or $J = R_1 \times L$ for some primary ideal L of R_2 . Hence, in view of Theorem 4, we have the following result.

Theorem 5. *Let $R = R_1 \times R_2$, where R_1 and R_2 are commutative rings with $1 \neq 0$, and let J be a proper ideal of R . The following statements are equivalent.*

- (1) J is a 1-absorbing primary ideal of R .
- (2) J is a primary ideal of R .
- (3) $J = I \times R_2$ for some primary ideal I of R_1 or $J = R_1 \times L$ for some primary ideal L of R_2 .

Recall that a nonzero nonunit element x of a ring R is called *irreducible* if $x = x_1x_2$ for some $x_1, x_2 \in R$, then x_1 is a unit of R or x_2 is a unit of R . Also, recall that x is called *prime* if $x \mid x_1x_2$ for some $x_1, x_2 \in R$, then $x \mid x_1$ or $x \mid x_2$. The following lemma is needed in the proof of our next result.

Lemma 2. *Let R be a quasilocal ring. If p is a nonzero prime element of R , then p is an irreducible element of R .*

Proof. Assume that $p = p_1p_2$ for some $p_1, p_2 \in R$. Since p is a prime element of R , we may assume that $p \mid p_1$. We show that p_2 is a unit of R . Hence $p_1 = pw$ for some $w \in R$ and thus $p = pwp_2$. Thus $p - pwp_2 = p(1 - wp_2) = 0$. If p_2 is a nonunit element of R , then $1 - wp_2$ is a unit of R (since R is quasilocal) and thus $p = 0$, a contradiction. Hence p_2 is a unit of R . Thus p is an irreducible element of R . \square

The following result provides a method to construct 1-absorbing primary ideals that are not primary ideals.

Theorem 6. *Let R be a quasilocal ring with maximal ideal M . Let $x \in M$ be a nonzero prime element of R such that $M \neq xR$. Then xM is a 1-absorbing primary ideal of R that is not a primary ideal of R .*

Proof. First, we show that xM is not a primary ideal of R . Observe that $\sqrt{xM} = xR$. Since $M \neq xR$, there exists an $m \in M \setminus xR$. Now $xm \in I$. Since x is a nonzero prime element of R , we conclude that x is an irreducible element of R by Lemma 2, and thus $x \notin xM$. Also, since $m \in M \setminus xR$ and $\sqrt{xM} = xR$, we conclude that $m \notin \sqrt{xM}$. Thus xM is not a primary ideal of R . Now we show that xM is a 1-absorbing primary ideal of R . Suppose that $abc \in I$ for some nonunit elements $a, b, c \in R$. Suppose that $ab \notin xM$. Then $x \nmid a$ and $x \nmid b$ (note that if $x \mid a$ or $x \mid b$, then $ab \in xM$). Since $x \mid abc$ and $x \nmid ab$, we conclude that $x \mid c$. Thus $c \in \sqrt{xM} = xR$. \square

Theorem 7. *Suppose that I is a 1-absorbing primary ideal of R that is not a primary ideal of R . Then there exist an irreducible element $x \in R$ and a nonunit element $y \in R$ such that $xy \in I$, but neither $x \in I$ nor $y \in \sqrt{I}$. Furthermore, if $ab \in I$ for some nonunit elements $a, b \in R$ such that neither $a \in I$ nor $b \in \sqrt{I}$, then a is an irreducible element of R .*

Proof. Since I is not a primary ideal of R , there exist nonunit elements $x, y \in R$ such that neither $x \in I$ nor $y \in \sqrt{I}$. Suppose that x is not an irreducible element of R . Then $x = cd$ for some nonunit elements $c, d \in R$. Since $xy = cdy \in I$ and I is a 1-absorbing primary ideal of R and $y \notin \sqrt{I}$, we conclude that $cd = x \in I$, a contradiction. Hence x is an irreducible element of R . \square

Theorem 8. *Let R be a quasilocal ring with maximal ideal M and P be a prime ideal of R such that $P \subseteq M$. Then PM is a 1-absorbing primary ideal of R .*

Proof. First observe that $\sqrt{PM} = P$. Suppose that $abc \in PM$ for some nonunit elements $a, b, c \in M$. If $a \in P$ or $b \in P$, then it is clear that $ab \in PM$. Hence assume that neither $a \in P$ nor $b \in P$. Thus $ab \notin P$. Since $abc \in PM \subseteq P$ and $ab \notin P$, we conclude that $c \in P = \sqrt{PM}$. Thus PM is a 1-absorbing primary ideal of R . \square

Remark 1. *Observe that PM in Theorem 8 needs not be a primary ideal of R by Theorem 6.*

Theorem 9. *Let I be a 1-absorbing primary ideal of a ring R . Suppose that $c \in R \setminus I$ is a nonunit element of R . Then $(I : c) = \{x \in R \mid cx \in I\}$ is a primary ideal of R .*

Proof. Suppose that I is a 1-absorbing primary ideal of R and c is a nonunit element of R such that $c \in R \setminus I$. Let $ab \in (I : c)$ for some elements $a, b \in R$. We may assume that a, b are nonunit elements of R . Suppose that $a \notin (I : c)$. Hence $ca \notin I$. Since $cab \in I$ and I is a 1-absorbing primary ideal of R and $ca \notin I$, we conclude that $b \in \sqrt{I} \subseteq \sqrt{(I : c)}$. Hence $(I : c)$ is a primary ideal of R . \square

Remark 2. *Let I be a 1-absorbing primary ideal of a ring R and c be a nonunit element of R such that $c \in R \setminus I$. Suppose that $c \notin \sqrt{I}$. Since \sqrt{I} is a prime ideal of R , we conclude that $I \subseteq (I : c) \subseteq \sqrt{I}$, and thus $\sqrt{(I : c)} = \sqrt{I}$. Suppose that $c \in \sqrt{I} \setminus I$. Let n be the last positive integer $n \geq 2$ such that $c^n \in I$. Then $c^{n-1} \in (I : c) \setminus I$. Thus $I \subsetneq (I : c)$. Also, $(I : c)$ needs not be a subset of \sqrt{I} ; for let R, I, M_M be as in Example 1, and $c = x$. Then $(I : c) = M_M \not\subseteq \sqrt{I}$.*

Recall that a ring R is called *divided* if for every prime ideal P of R and for every $x \in R \setminus P$, we have $x \mid p$ for every $p \in P$. Recall that a ring R is called a *chained* ring if for every $x, y \in R$, we have $x \mid y$ or $y \mid x$. Thus every chained ring is divided. Hence if R is a divided ring, then R is a quasilocal ring. We have the following result.

Theorem 10. *Let R be a divided ring. Then a proper ideal I of R is a 1-absorbing primary ideal of R if and only if I is a primary ideal of R . In particular, if R is a chained ring, then a proper ideal I of R is a 1-absorbing primary ideal of R if and only if I is a primary ideal of R .*

Proof. It is clear that every primary ideal of R is a 1-absorbing primary ideal of R . Hence assume that I is a 1-absorbing primary ideal of R . Suppose that $xy \in I$ for some $x, y \in R$ and $y \notin \sqrt{I}$. We may assume that x, y are nonunit elements of R . Since \sqrt{I} is a prime ideal of R by Theorem 2 and $y \notin \sqrt{I}$, we conclude that $x \in \sqrt{I}$. Since R is divided, we conclude that $y \mid x$. Thus $x = yw$ for some $w \in R$. Since $y \notin \sqrt{I}$ and $x \in \sqrt{I}$, we conclude that w is a nonunit element of R . Since $xy = ywy \in I$ and I is a 1-absorbing primary ideal of R and $y \notin \sqrt{I}$, we conclude that $yw = x \in I$. Thus I is a primary ideal of R . \square

Recall that a proper ideal I of R is called *principal* if $I = xR$ for some $x \in R$.

Theorem 11. *Let R be a divided ring with maximal ideal M . If M is not a principal prime ideal of R , then every nonzero prime ideal of R is not principal.*

Proof. Suppose that a nonzero prime ideal $P = xR$ for some $x \in R$. Then x is a nonzero prime element of R and $M \neq xR$. Thus xM is a 1-absorbing primary ideal of R that is not a primary ideal of R by Theorem 6, which is a contradiction by Theorem 10. \square

Theorem 12. *Let R be a divided integral domain and P be a prime ideal of R . Then P^n is a primary ideal of R for every positive integer $n \geq 1$, and hence P^n is a 1-absorbing primary ideal of R for every positive integer $n \geq 1$.*

Proof. If $n = 1$, then there is nothing to prove. Thus let $n \geq 2$ and Suppose that $xy \in P^n$ for some $x, y \in R$. Then $xy = p_1c_1 + \cdots + p_kc_k \in P^n$ for some $p_1, \dots, p_k \in P$ and $c_1, \dots, c_k \in P^{n-1}$ for some positive integer $k \geq 1$. Suppose that $y \notin P$. Since R is divided, we have Then $y \mid p_i$ for every i , $1 \leq i \leq k$. Hence for every i , $1 \leq i \leq k$, we have $p_i = yd_i$ for some $d_i \in P$. Thus $xy = yd_1c_1 + \cdots + yd_kc_k$. Hence $y(x - (d_1c_1 + \cdots + d_kc_k)) = 0$. Since R is an integral domain, we conclude that $x = d_1c_1 + \cdots + d_kc_k \in P^n$. Thus P^n is a primary ideal of R . Since every primary ideal of R is a 1-absorbing primary ideal, we conclude that P^n is a 1-absorbing primary ideal of R for every positive integer $n \geq 2$. \square

Recall that an integral domain R is called a *valuation domain* if R is a chained ring.

Theorem 13. *Let R be a valuation domain and I be a proper ideal of R with $\sqrt{I} = P$ (note that P is a prime ideal of R). The following statements are equivalent.*

- (1) I is a 1-absorbing primary ideal of R .
- (2) I is a primary ideal of R .
- (3) If $P \neq P^2$, then $I = P^n$ for some positive integer $n \geq 1$.

Proof. (1) \Rightarrow (2). Since R is divided, the claim is clear by Theorem 10.

(2) \Rightarrow (3). The claim is clear by [8, Theorem 5.11].

(3) \Rightarrow (1). Since R is divided, the claim is clear by Theorem 12. \square

Let R be an integral domain with quotient field K . Recall that a proper ideal I of R is called *invertible* if $II^{-1} = R$, where $I^{-1} = \{r \in K \mid rI \subseteq R\}$. An integral domain R is called a *Prufer domain* if every nonzero finitely generated ideal of R is invertible.

Theorem 14. *Let R be a Prufer domain and I be a proper ideal of R with $\sqrt{I} = P$ for some prime ideal P of R . Then the following statements are equivalent.*

- (1) I is a 1-absorbing primary ideal of R .
- (2) I is a primary ideal of R .
- (3) If P is a finitely generated ideal of R , then $I = P^n$ for some positive integer $n \geq 1$.

Proof. (1) \Rightarrow (2). Suppose that R is quasilocal with maximal ideal M . Since R is a Prufer domain, it is known that $R = R_M$ is a valuation domain and hence the claim follows from Theorem 14. Suppose that R is not quasilocal. Then the claim follows by Theorem 4.

(2) \Rightarrow (3). The claim is clear by [8, Exercise 2, p. 144].

(3) \Rightarrow (1). Since $I = P^n$ for some positive integer $n \geq 1$ and P is a finitely generated ideal of R , we conclude that I is a primary ideal of R by [8, Exercise 2, p. 144]. Thus I is a 1-absorbing primary ideal of R . \square

Recall that an integral domain is called a *Dedekind domain* if every nonzero proper ideal of R is invertible.

Theorem 15. *Let R be a Dedekind domain and I be a nonzero proper ideal of R . Then I is a 1-absorbing primary ideal of R if and only if \sqrt{I} is a prime ideal of R .*

Proof. If I is a 1-absorbing primary ideal of R , then \sqrt{I} is a prime ideal of R by Theorem 2. Conversely, suppose \sqrt{I} is a prime ideal of R . Since R is a Dedekind domain, it is well-known that every nonzero prime ideal of R is a maximal ideal of R . Thus \sqrt{I} is a maximal ideal of R . Hence I is a primary ideal of R , and thus I is 1-absorbing primary ideal of R . \square

Theorem 16. *Let R be a Noetherian integral domain that is not a field. Then the following statements are equivalent:*

- (1) R is a Dedekind domain.
- (2) A nonzero proper ideal I of R is a 1-absorbing primary ideal of R if and only if $I = P^n$ for some prime ideal P of R some positive integer n .

Proof. (1) \Rightarrow (2). Let R be a Dedekind domain and I be a nonzero proper ideal of R . Suppose that I is a 1-absorbing primary ideal of R . Then $\sqrt{I} = P$ is a nonzero prime ideal of R by Theorem 2. Since R is a Dedekind domain, it is known that every nonzero prime ideal of R is a maximal ideal of R . Thus $\sqrt{I} = P$ is a maximal ideal of R . Thus I is a primary ideal of R . Since R is Dedekind and I is a primary ideal of R with $\sqrt{I} = P$ is a maximal ideal of R , we conclude that $I = P^n$ for some $n \geq 1$ by of R by [8, Theorem 6.20]. Conversely, suppose that $I = P^n$ for some nonzero proper ideal P of R and a positive integer $n \geq 1$. Since R is Dedekind, we conclude that P is a maximal ideal of R , and hence I is a primary ideal of R . Thus I is a 1-absorbing primary ideal of R .

(2) \Rightarrow (1). Suppose that every nonzero 1-absorbing primary ideals of R is of type $I = P^n$ for some nonzero prime ideal P of R and a positive integer $n \geq 1$. Let M be a maximal ideal of R . Since every ideal between M^2 and M is a primary ideal and hence a 1-absorbing primary ideal of R , we conclude that there is no primary ideals of R between M^2 and M . Thus R is a Dedekind domain by [5, Theorem 39.2]. \square

Since every principal ideal domain is a Dedekind domain, we have the following result.

Corollary 1. *Let R be a principal ideal domain and I be a nonzero proper ideal of R . Then I is a 1-absorbing primary ideal of R if and only if $I = p^n R$ for some nonzero prime element p of R and a positive integer $n \geq 1$.*

In light of Theorem 6 and Example 1, observe that there are some rings whose 1-absorbing primary ideals are not of the form P^n for some prime ideal P of R and a positive integer $n \geq 1$.

Let I_1 and I_2 be 1-absorbing primary ideals of R . If $\sqrt{I_1} \neq \sqrt{I_2}$, then $I_1 \cap I_2$ needs not to be a 1-absorbing primary ideal of R . We have the following example.

Example 3. *Let $R = \mathbb{Z} \times \mathbb{Z}$. Then $I_1 = 4\mathbb{Z} \times \mathbb{Z}$ and $I_2 = \mathbb{Z} \times 9\mathbb{Z}$ are 1-absorbing primary ideals of R . Also, $\sqrt{I_1} = 2\mathbb{Z} \times \mathbb{Z}$ and $\sqrt{I_2} = \mathbb{Z} \times 3\mathbb{Z}$. Hence $\sqrt{I_1} \neq \sqrt{I_2}$ and $I_1 \cap I_2 = 4\mathbb{Z} \times 9\mathbb{Z}$ is not a 1-absorbing primary ideal of R by Theorem 5*

Definition 2. Let I be a 1-absorbing primary ideal of a ring R . Then $\sqrt{I} = P$ is a prime ideal of R by Theorem 2. Hence we call I a P -1-absorbing primary ideal of R .

So we have the following result.

Theorem 17. Let I_1, I_2, \dots, I_n be P -1-absorbing primary ideals of a ring R . Then $I = \bigcap_{i=1}^n I_i$ is a P -1-absorbing primary ideal of R .

Proof. First observe that $\sqrt{I} = P$. Suppose that $abc \in I$ for some nonunit elements $a, b, c \in R$ and $ab \notin I$. Without loss of generality, we may assume that $ab \notin I_1$. Since I_1 is a P -1-absorbing primary ideal of R and $ab \notin I_1$, we have $c \in P$. \square

Theorem 18. Let R_1 and R_2 be rings and $f : R_1 \rightarrow R_2$ be a ring homomorphism such that $f(1) = 1$ and if R_2 is a quasilocal ring, then $f(a)$ is a nonunit of R_2 for every nonunit $a \in R_1$. Then the following statements hold.

- (1) Assume that J is a 1-absorbing primary ideal of R_2 , then $f^{-1}(J)$ is a 1-absorbing primary ideal of R_1 .
- (2) If f is onto and I is a 1-absorbing primary ideal of R_1 with $\text{Ker}(f) \subseteq I$, then $f(I)$ is a 1-absorbing primary ideal of R_2 .

Proof. (1). Assume that R_2 is quasilocal. Let $abc \in f^{-1}(J)$ for some nonunit elements $a, b, c \in R_1$. Then $f(abc) = f(a)f(b)f(c) \in J$ (note that $f(a), f(b)$, and $f(c)$ are nonunit elements of R_2 by hypothesis), which implies $f(a)f(b) \in J$ or $f(c) \in \sqrt{J}$. It follows $ab \in f^{-1}(J)$ or $c \in \sqrt{f^{-1}(J)} = f^{-1}(\sqrt{J})$. Thus $f^{-1}(J)$ is a 1-absorbing primary ideal of R_1 . Suppose that R_2 is not a quasilocal ring and J is a 1-absorbing primary ideal of R_2 . Then J is a primary ideal of R_2 by Theorem 3. Thus $f^{-1}(J)$ is a primary ideal of R_1 , and hence $f^{-1}(J)$ is a 1-absorbing primary ideal of R_1 .

(2). Since f is onto and $\text{Ker}(f) \subseteq I$, we know that $f(\sqrt{I}) = \sqrt{f(I)}$. Let $xyz \in f(I)$ for some nonunit elements $x, y, z \in R_2$. Since f is onto, there exist nonunit elements $a, b, c \in R_1$ such that $x = f(a)$, $y = f(b)$, and $z = f(c)$. Hence $f(abc) = f(a)f(b)f(c) = xyz \in f(I)$. Since $\text{Ker}(f) \subseteq I$, we conclude that $abc \in I$. Hence $ab \in I$ or $c \in \sqrt{I}$; so $xy \in f(I)$ or $z \in f(\sqrt{I}) = \sqrt{f(I)}$. Thus $f(I)$ is a 1-absorbing primary ideal of R_2 . \square

In view of Theorem 18, we have the following result.

Corollary 2. Let I and J be proper ideals of a ring R with $I \subseteq J$ and suppose that if R/I is a quasilocal ring, then $a + I$ is a nonunit of R/I for every nonunit $a \in R$. Then J is a 1-absorbing primary ideal of R if and only if J/I is a 1-absorbing primary ideal of R/I .

Proof. Let $f : R \rightarrow R/I$ such that $f(a) = a + I$. Then f is a ring homomorphism from R onto R/I and $f(1) = 1$. Suppose that J is a 1-absorbing primary ideal of R . Since $\text{Ker}(f) = I \subseteq J$ and f is onto, we conclude that $f(J) = J/I$ is a 1-absorbing primary ideal of R/I by Theorem 18(2). Suppose that J/I is a 1-absorbing primary ideal of R/I . Then $f^{-1}(J/I) = J$ is a 1-absorbing primary ideal of R by Theorem 18(1). \square

Theorem 19. Let S be a multiplicatively closed subset of R , and I be a proper ideal of R . Then the following statements hold.

- (1) If I is a 1-absorbing primary ideal of R such that $I \cap S = \emptyset$, then $S^{-1}I$ is a 1-absorbing primary ideal of $S^{-1}R$.
- (2) If $S^{-1}I$ is a 1-absorbing primary ideal of $S^{-1}R$ and $S \cap Z_I(R) = \emptyset$, then I is a 1-absorbing primary ideal of R .

Proof. (1) Let $\frac{a}{s_1} \frac{b}{s_2} \frac{c}{s_3} \in S^{-1}I$ for some nonunit elements $a, b, c \in R$, and $s_1, s_2, s_3 \in S$. Suppose that $\frac{a}{s_1} \frac{b}{s_2} \notin S^{-1}I$. Then $uabc \in I$ for some $u \in S$. Since I is 1-absorbing primary and $uab \notin I$, we have $c \in \sqrt{I}$. Thus $\frac{c}{s_3} \in S^{-1}\sqrt{I} = \sqrt{S^{-1}I}$ which completes the proof.

(2) Let $abc \in I$ for some nonunit elements $a, b, c \in R$. Hence $\frac{abc}{1} = \frac{a}{1} \frac{b}{1} \frac{c}{1} \in S^{-1}I$. Since $S^{-1}I$ is 1-absorbing primary, we have $\frac{a}{1} \frac{b}{1} \in S^{-1}I$ or $\frac{c}{1} \in \sqrt{S^{-1}I} = S^{-1}\sqrt{I}$. If $\frac{a}{1} \frac{b}{1} \in S^{-1}I$, then $uab \in I$ for some $u \in S$. Since $u \notin Z_I(R)$, we conclude that $ab \in I$. If $\frac{c}{1} \in S^{-1}\sqrt{I}$, then $(tc)^n \in I$ for some positive integer n and $t \in S$. Since $t^n \notin Z_I(R)$, we have $c^n \in I$, as needed. \square

Theorem 20. *Let I be a 1-absorbing primary ideal of a commutative ring R . If $abJ \subseteq I$ for some nonunit elements $a, b \in R$ and a proper ideal J of R , then $ab \in I$ or $J \subseteq \sqrt{I}$.*

Proof. Assume on the contrary that $abJ \subseteq I$, but $ab \notin I$ and $J \not\subseteq \sqrt{I}$. Then there exists an element $j \in J$ such that $j \notin \sqrt{I}$. Hence we have $abj \in I$, but neither $ab \in I$ nor $j \in \sqrt{I}$, a contradiction. \square

Theorem 21. *Let I be a proper ideal of R . Then the followings statements are equivalent.*

- (1) I is a 1-absorbing primary ideal of R .
- (2) For any proper ideals I_1, I_2, I_3 of R such that $I_1I_2I_3 \subseteq I$ implies that either $I_1I_2 \subseteq I$ or $I_3 \subseteq \sqrt{I}$.

Proof. (1) \Rightarrow (2). Let I be a 1-absorbing primary ideal of R . Assume that $I_1I_2I_3 \subseteq I$ for some proper ideals I_1, I_2, I_3 of R and $I_1I_2 \not\subseteq I$. Then there exist nonunit elements $a \in I_1$ and $b \in I_2$ such that $ab \notin I$. Since $abI_3 \subseteq I$ and $ab \notin I$, we conclude that $I_3 \subseteq \sqrt{I}$ by Theorem 20.

(2) \Rightarrow (1). Suppose that $abc \in I$ for some nonunit elements $a, b, c \in R$ and $ab \notin I$. Let $I_1 = aR, I_2 = bR$, and $I_3 = cR$. Then $I_1I_2I_3 \subseteq I$ and $I_1I_2 \not\subseteq I$. Thus $I_3 = cR \subseteq \sqrt{I}$. Hence $c \in \sqrt{I}$. \square

REFERENCES

- [1] D. D. Anderson and M. Batanieh, Generalizations of prime ideals, *Comm. Algebra*, **36** (2008), 686–696.
- [2] D.F. Anderson, A. Badawi, On n -absorbing ideals of commutative rings, *Comm. Algebra*, **39** (2011), 1646–1672.
- [3] A. Badawi, On 2-absorbing ideals of commutative rings, *Bull. Austral. Math. Soc.*, **75** (2007), 417–429.
- [4] A. Badawi, Ü. Tekir and E. Yetkin, On 2-absorbing primary ideals in commutative rings, *Bull. Korean Math. Soc.*, **51** (4) (2014), 1163–1173.
- [5] R. Gilmer, *Multiplicative ideal theory*, Queen Papers Pure Appl. Math. 90, Queen’s University, Kingston, 1992.
- [6] J. Huckaba, *Rings with zero-divisors*, Marcel Dekker, NewYork/ Basil, 1988.
- [7] I. Kaplansky, *Commutative rings*, rev. ed., University of Chicago Press, Chicago, 1974.
- [8] M.D. Larson and P.J. McCarthy, *Multiplicative theory of ideals*, Academic Press, New York, London, 1971.

DEPARTMENT OF MATHEMATICS & STATISTICS, THE AMERICAN UNIVERSITY OF SHARJAH, P.O.
BOX 26666, SHARJAH, UNITED ARAB EMIRATES.

Email address: `abadawi@aus.edu`

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE AND ART, GAZIANTEP UNIVERSITY,
GAZIANTEP, TURKEY

Email address: `yetkinece@gmail.com`