Stability and bifurcation analysis of a discrete-time predator-prey model with strong Allee effect

Ziyad AlSharawi[†], Saheb Pal^a, Sourav Rana^b, Nikhil Pal^a, Joydev Chattopadhyay^{c1}

[†] *Department of Mathematics and Statistics, American University of Sharjah, UAE*

^a *Department of Mathematics, Visva-Bharati, Santiniketan 731235, India*

^b *Department of Statistics, Visva-Bharati, Santiniketan 731235, India*

^c *Agricultural and Ecological Research Unit, Indian Statistical Institute
203, B. T. Road, Kolkata 700108, India*

Abstract

In this paper, we investigate the impact of Allee effect on the stability of a discrete-time predator-prey model with a non-monotonic functional response. The proposed model supports the coexistence of two steady states, and the mathematical features of the model are analyzed based on local stability and bifurcation theory. By considering the Allee parameter as the bifurcation parameter, we provide sufficient conditions for the period-doubling and the Neimark-Sacker bifurcations. Also, sufficient conditions are given to show that increasing the magnitude of the Allee parameter within a certain domain plays a significant role in stabilizing the system.

Keywords: Predator-prey model; Allee effect; Local stability; Period-doubling bifurcation; Neimark-Sacker bifurcation.

1 Introduction

There are various modelling approaches in the fields of mathematical biology and ecology, and among which discrete-time and continuous-time modelling. Discrete-time mathematical models are widely used in population ecology [1], which can be attributed to several factors; among which: population growth at discrete intervals of time, numerical simulations of non-discrete models, and the capability of simple discrete models to cover a wide range of complex behaviour. Firstly, when population growth takes place at discrete units of time and generations are non-overlapping, discrete-time models are the appropriate mathematical tool to use [2]. Secondly, it is quite common to simulate nonlinear continuous models through certain type of discretization [3], which triggers the issue of starting with a discrete model in the first place. Finally, discrete-time models exhibit rich dynamics compared to continuous-time models. For instance, chaos can be observed in a single-species discrete model [1], whereas at least three species model is needed for continuous autonomous systems [4].

After the pioneering work of Lotka [5] and Volterra [6], two-dimensional spatially homogeneous models of predator-prey interactions started to attract the interest of many researchers. For modelling the predator-prey interactions, the growth of the prey population to its carrying capacity in the absence of predators is a main issue that needs to be considered. For instance, with logistic growth. This logistic growth assumes per capita growth rate of prey to be maximum when their

¹ *Corresponding author* E-mail: joydev@isical.ac.in.

population density is low and starts decaying with the increase in prey density. But, such growth rate is not always positive for small density [7; 8]. There are several biological reasons behind this. For example at low density, individuals will not always be able to find a suitable mate.

In 1931, an ecologist Warder Clyde Allee described that prey growth rate is negative or an increasing function at low population density [7]. This type of mechanism is known as Allee effect [9; 10]. There are several biological and environmental factors behind this Allee effect, such as reproductive facilitation, complications in finding mates, predation, environment conditioning, inbreeding depression, etc. [7; 8]. In short, a population is said to have an Allee effect, if there is a positive relation between the population density and the per capita growth rate at low density. Empirical evidence of Allee effects has been observed in many natural species, for instance, plants [11; 12], insects [13], marine invertebrates [14], birds and mammals [15]. Allee effects can be mainly classified into two broad categories: strong and weak [8]. There is a threshold population level for the strong Allee effect such that the species become extinct below the threshold population density. On the other hand, the weak Allee effect occurs when the growth rate reduces but remains positive at low population density [16; 17; 18; 19]. Many researchers have paid attention to study the dynamics of predator-prey models with Allee effect in prey growth rate [20; 21; 22].

To incorporate Allee phenomenon in the predator-prey model, there is a dominant formalism to modify the logistic growth rate of prey population. The relevant functional form is given by

$$f(x) = r \left(1 - \frac{x}{k}\right) (x - \theta)$$

and is known as “prey growth rate with multiplicative Allee effect” [9; 10]. Here x is the population density, $r > 0$ is the intrinsic growth rate and $k > 0$ is the carrying capacity of the prey population. The Allee parameter θ satisfies $|\theta| < k$. The Allee effect is said to be strong if $0 < \theta < k$. For the strong Allee effect, the function $f(x)$ satisfies the following properties: (i) $f(x)$ is negative for $0 < x < \theta$ and $x > \theta$, positive for $\theta < x < k$ (ii) $f'(x)$ is positive for $0 < x < (k + \theta)/2$, negative for $x > (k + \theta)/2$ and attains the maximum at $x = (k + \theta)/2$. Other types of mathematical formulation to model the Allee effect in predator-prey interactions are given in [8].

To describe the predator-prey interaction one key element that reflects the behavioral characteristic of the predator species is called the functional response or trophic function. Eventually, the functional response plays a dominant role in determining the different complex dynamical behaviors, namely, the steady states, bistability, periodic oscillations, chaos and bifurcation phenomena [23]. An inherent assumption that is true for many predator-prey interactions is that the functional response is monotonic. However, many observational and experimental studies propose that it may be non-monotonic, for example in the cases of “inhibition” in microbial dynamics and “group defense” in population dynamics [24; 25]. The term “group defense” used to describe the phenomenon when prey populations are large enough, predation process is decreased due to increased the ability of prey to better defend. In the present paper, we consider a simplified Monod-Haldane functional form $p(x) = mx/(x^2 + a)$ as a non-monotonic functional response, which is also known as Holling type-IV functional response.

In general, it has been shown that replacing the logistic growth function with Allee-type function could dramatically change the patterns of the dynamics, thus there is a need for mathematical analysis of predator-prey models with more realistic Allee effect growth functions. The present paper deals with a discrete-time predator-prey model subject to strong Allee effect on prey population and predator population predate the prey population in a simplified Monod-Haldane functional response. The rest of the paper is organized as follows: In Section 2, we present the mathematical model with its assumptions and steady states. In Section 3, we characterize the local stability of the equilibrium solutions. In Section 4, we study the period-doubling as well as Neimark-Sacker bifurcations for our proposed model with respect to the Allee parameter. Finally, we close this paper with a brief conclusion.

2 The model and its equilibria

We consider the discrete-time predator-prey model

$$\begin{cases} x_{n+1} = x_n \exp \left[r \left(1 - \frac{x_n}{k} \right) (x_n - \theta) - \frac{y_n}{x_n^2 + a} \right] =: x_n F(x_n, y_n) \\ y_{n+1} = y_n \exp \left[\frac{\mu x_n}{a + x_n^2} - d \right] =: y_n G(x_n), \end{cases} \quad (2.1)$$

where x_n and y_n are the density of the prey and predator, respectively. n is a non-negative integer, $r > 0$ is the intrinsic growth rate, μ is the conversion coefficient of prey biomass to predator biomass lies between 0 and 1, $d > 0$ is the death rate of the predator, a is a positive parameter. It is natural here to assume that $0 < \theta < k$. This model is a discrete version of the continuous-time predator-prey model with strong Allee effect

$$\begin{cases} \frac{dx}{dt} = rx \left(1 - \frac{x}{k} \right) (x - \theta) - \frac{xy}{x^2 + a}, \\ \frac{dy}{dt} = \frac{\mu xy}{x^2 + a} - dy, \end{cases} \quad (2.2)$$

where the predator prey interaction follows a non-monotonic functional response. Indeed, a known nonstandard discretization method is to consider piecewise constant arguments [26; 3] as follows:

$$\begin{aligned} \frac{1}{x(t)} \frac{dx(t)}{dt} &= r \left(1 - \frac{x([t])}{k} \right) (x([t]) - \theta) - \frac{y([t])}{x^2([t]) + a}, \\ \frac{1}{y(t)} \frac{dy(t)}{dt} &= \frac{\mu x([t])}{x^2([t]) + a} - d, \end{aligned} \quad (2.3)$$

where $0 \leq n \leq t < n+1$ and $[t]$ is the greatest integer in t . Now, observe that the right hand side of the system is constant over the interval $[n, n+1)$. Thus, by integration over the interval $[n, n+1)$, we obtain

$$\begin{aligned} x(n+1) &= x(n) \exp \left[r \left(1 - \frac{x(n)}{k} \right) (x(n) - \theta) - \frac{y(n)}{x^2(n) + a} \right], \\ y(n+1) &= y(n) \exp \left[\frac{\mu x(n)}{x^2(n) + a} - d \right], \end{aligned} \quad (2.4)$$

for any $t \in [n, n+1)$, $n = 0, 1, 2, \dots$. Finally, let $x(n) = x_n$ and $y(n) = y_n$ to obtain the discrete-time model in System (2.1). Obviously, the initial values $x(0) = x_0$ and $y(0) = y_0$ are assumed to

be non-negative real numbers.

In the absence of the predator, the model in Eq. (2.1) reduces to the one dimensional Ricker model with Allee effects, namely

$$x_{n+1} = x_n e^{r(1-\frac{x_n}{k})(x_n-\theta)} =: R(x_n), \quad 0 < \theta < k. \quad (2.5)$$

The Allee effects in one dimensional discrete-time models have been investigated in the literature, and the interested reader can refer to [27; 28] for more details. In particular, Eq. (2.5) has the equilibrium solutions $\bar{x} = 0, \theta$ and k . The equilibrium $\bar{x} = 0$ is always stable while $\bar{x} = \theta$ is always unstable. On the other hand, $\bar{x} = k$ is stable when $r(k - \theta) < 2$ and loses its stability when $r(k - \theta) > 2$. Observe that the equation $R(x) = \theta$ has two solutions, one of them is θ and the other one (say α_θ) is larger than k . Initial populations that are less than θ or larger than α_θ lead to a collapse of the population. This phenomenon carries over to the predator-prey model in System (2.1), which naturally leads to the collapse in both species.

Next, we focus on the equilibrium solutions of System (2.1) when both species exist. Thus, we solve the system of equations

$$\begin{cases} x = xF(x, y) = x \exp \left[r(1 - \frac{x}{k})(x - \theta) - \frac{y}{x^2 + a} \right] \\ y = yG(x) = y \exp \left[\frac{\mu x}{x^2 + a} - d \right]. \end{cases} \quad (2.6)$$

Obviously, $E_0 := (0, 0)$, $E_{\bar{x}_1} := (\theta, 0)$ and $E_{\bar{x}_2} := (k, 0)$ are three equilibrium solutions that we call the trivial equilibria. To investigate the existence of nontrivial ones, assume $xy \neq 0$. When $\mu^2 - 4ad^2 > 0$, we obtain two solutions of the equation $1 = G(x)$ that merge at $\mu^2 - 4ad^2 = 0$. Because $\mu^2 - 4ad^2 \leq 0$ implies $\frac{\mu x_n}{a + x_n^2} - d \leq 0$, and consequently, $y_{n+1} \leq y_n$, then the predator is doomed to extinction. Therefore, it is natural to proceed under the assumption that $\mu^2 - 4ad^2 > 0$. In this case, we need the two positive solutions of $G(x) = 1$ to satisfy the equation $1 = F(x, y)$. Otherwise, System (2.1) has no nontrivial positive equilibrium solutions, which makes this scenario beyond our interest. Denote the positive solutions of $G(x) = 1$ by

$$\bar{x}_4 := \frac{\mu}{2d} - \sqrt{\left(\frac{\mu}{2d}\right)^2 - a} \quad \text{and} \quad \bar{x}_5 := \frac{\mu}{2d} + \sqrt{\left(\frac{\mu}{2d}\right)^2 - a},$$

then the conditions $\theta < \bar{x}_4$ and $\bar{x}_5 < k$ guarantee the existence of two nontrivial equilibrium solutions (coexistence solutions), namely

$$E_4 := (\bar{x}_4, \bar{y}_4) \text{ and } E_5 := (\bar{x}_5, \bar{y}_5), \quad \text{where} \quad \bar{y}_j = r(\bar{x}_j^2 + a)(1 - \frac{\bar{x}_j}{k})(\bar{x}_j - \theta) \text{ for } j = 4, 5.$$

Up to this end, we have established the existence of all equilibrium solutions under the assumption

$$a < \left(\frac{\mu}{2d}\right)^2. \quad (2.7)$$

In general, it is possible to have no coexistence equilibrium, one coexistence equilibrium or two coexistence equilibria. Figure 1 gives an illustration.

Figure 1: This figure illustrates the trivial and nontrivial equilibrium solutions as discussed above. The parameters are taken to be $a = 6, d = 1, \mu = 4, \theta = 1, k = 4$ and $r = \frac{1}{5}$. The red curve is $F(x, y) = 1$, which is the nullcline (or zero isocline) of the prey equation while the blue vertical lines are the nullcline of the predator equation, i.e., $G(x) = 1$. The small arrows indicate orbits direction in each region.

3 Local stability analysis

Consider the assumption in inequality (2.7). In this section, we investigate the local stability of the equilibrium solutions in three cases, namely $\theta = \bar{x}_4 < \bar{x}_5 < k$, $\theta < \bar{x}_4 < \bar{x}_5 = k$ and $\theta < \bar{x}_4 < \bar{x}_5 < k$. We stress those cases in the following proposition.

Proposition 3.1. *Assume that $\mu > 2d\sqrt{a}$. Each of the following holds true:*

(i) $\theta = \bar{x}_4 < \bar{x}_5 < k$ if and only if

$$\frac{\mu}{d} = \theta + \frac{a}{\theta} \quad \text{and} \quad \frac{\mu}{d} < k + \frac{a}{d}.$$

(ii) $\theta < \bar{x}_4 < \bar{x}_5 = k$ if and only if

$$\frac{\mu}{d} < \theta + \frac{a}{\theta} \quad \text{and} \quad \frac{\mu}{d} = k + \frac{a}{k}.$$

(iii) $\theta < \bar{x}_4 < \bar{x}_5 < k$ if and only if

$$\frac{\mu}{d} < \min \left\{ \theta + \frac{a}{\theta}, k + \frac{a}{k} \right\}.$$

Thus, we have either one or two coexistence equilibrium, and that will be the scenarios we consider in our stability analysis. The Jacobian matrix of System (2.1) at an equilibrium solution $(\bar{x}, \bar{y})$ is given by

$$J(\bar{x}, \bar{y}) = \begin{bmatrix} \left(1 + r\bar{x} \left(1 + \frac{\theta}{k} - \frac{2\bar{x}}{k} \right) + \frac{2\bar{x}^2\bar{y}}{(\bar{x}^2+a)^2} \right) F(\bar{x}, \bar{y}) & \frac{-\bar{x}}{\bar{x}^2+a} F(\bar{x}, \bar{y}) \\ \bar{y} G(\bar{x}) \frac{\mu(a-\bar{x}^2)}{(\bar{x}^2+a)^2} & G(\bar{x}) \end{bmatrix}. \quad (3.1)$$

At a nontrivial equilibrium $(\bar{x}, \bar{y})$, we have $G(\bar{x}) = 1$ and $F(\bar{x}, \bar{y}) = 1$, and consequently,

$$\frac{\mu\bar{x}}{\bar{x}^2+a} = d \quad \text{and} \quad \frac{\bar{y}}{\bar{x}^2+a} = r \left(1 - \frac{\bar{x}}{k} \right) (\bar{x} - \theta). \quad (3.2)$$

Thus, the Jacobian matrix in Eq. (3.1) becomes

$$J(\bar{x}, \bar{y}) = \begin{bmatrix} \left(1 + r\bar{x} \left(1 + \frac{\theta}{k} - \frac{2\bar{x}}{k}\right) + 2\frac{\bar{x}^2\bar{y}}{(\bar{x}^2+a)^2}\right) & \frac{-d}{\mu} \\ \bar{y}\frac{\mu(a-\bar{x}^2)}{(\bar{x}^2+a)^2} & 1 \end{bmatrix}. \quad (3.3)$$

The trace and determinant of J are given by

$$\begin{aligned} T := \text{Trace}(J) &= 2 + \frac{2r}{k}\bar{x} \left(\frac{k+\theta}{2} - \bar{x}\right) + \frac{2\bar{x}^2\bar{y}}{(\bar{x}^2+a)^2} \quad \text{and} \\ D := \text{Det}(J) &= T - 1 + \bar{y}\frac{d(a-\bar{x}^2)}{(\bar{x}^2+a)^2} \\ &= 1 + \frac{2r}{k}\bar{x} \left(\frac{k+\theta}{2} - \bar{x}\right) + \frac{\bar{y}}{(\bar{x}^2+a)^2} (2\bar{x}^2 + d(a-\bar{x}^2)). \end{aligned}$$

Recall that if both eigenvalues of J are within the unit disk, then the equilibrium is locally asymptotically stable and it is called a sink. If both eigenvalues are out of the unit disk, the equilibrium is unstable, and it is called a source. If one eigenvalue is within and the other one is outside of the unit disk, then it is called a saddle. If the eigenvalues of J do not satisfy any of the aforementioned scenarios, then the equilibrium is called non-hyperbolic, and in this case, investigating local stability needs nonstandard techniques. Denote the trace and determinant of the matrix in Eq. (3.1) by T and D , respectively. The characteristic equation of the Jacobian is given by $P(\lambda) = \lambda^2 - T\lambda + D$. Here, it is convenient to characterize local stability based on the trace-determinant values. Write the eigenvalues as

$$\lambda_1 = \frac{1}{2} \left(T - \sqrt{T^2 - 4D} \right) \quad \text{and} \quad \lambda_2 = \frac{1}{2} \left(T + \sqrt{T^2 - 4D} \right),$$

then the graph in Fig. 2 facilitates our stability analysis, which follows based on the above cases.

Figure 2: This figure illustrates the magnitude of the eigenvalues based on the trace-determinant values. T is used for trace and D is used for determinant. Also, $\|\cdot\|$ is used for complex eigenvalues, whereas $|\cdot|$ is used for real eigenvalues.

The next classical lemma characterizes stability based on the trace-determinant values.

Lemma 3.2. *Let $(\bar{x}, \bar{y})$ be an equilibrium solution of System (2.1). If T is the trace and D is the determinant of the Jacobian matrix at $(\bar{x}, \bar{y})$, then each of the following holds true:*

(i) $(\bar{x}, \bar{y})$ is a sink if

$$-1 < D < 1 \quad \text{and} \quad -D - 1 < T < D + 1.$$

(ii) $(\bar{x}, \bar{y})$ is a source if

$$D > 1 \quad \text{and} \quad -D - 1 < T < D + 1, \quad \text{or} \quad D < -1 \quad \text{and} \quad D + 1 < T < -D - 1.$$

(iii) $(\bar{x}, \bar{y})$ is a saddle if

$$0 < |T| + D + 1 < 2|T|.$$

(iv) $(\bar{x}, \bar{y})$ is nonhyperbolic if

$$|T| = |D + 1|, \quad \text{or} \quad D = 1 \quad \text{and} \quad |T| \leq 2.$$

Before we proceed into particular cases, we give the next proposition, which shows how E_4 and E_5 fit within Lemma 3.2.

Proposition 3.3. *Consider $\mu^2 > 4ad^2$. Each of the following holds true:*

(i) $\bar{x}_4 < \sqrt{a} < \bar{x}_5$.

(ii) $D > T - 1$ at E_4 , and $D < T - 1$ at E_5 .

Proof. (i) Recall that $\bar{x}_4$ and $\bar{x}_5$ are the zeros of $p(x) = dx^2 - \mu x + ad$. They show up at $\mu^2 = 4ad^2$, in which $\bar{x}_4 = \bar{x}_5 = \sqrt{a}$ and consequently, $D + 1 - T = 0$. When $\mu^2 > 4ad^2$, we have $\bar{x}_5 > \bar{x}_4$ and $\bar{x}_4 \bar{x}_5 = a$. Thus, $\bar{x}_4 < \sqrt{a} < \bar{x}_5$. For Part (ii), we have

$$D - T + 1 = d\bar{y} \frac{a - \bar{x}^2}{(a + \bar{x}^2)^2}, \quad (3.4)$$

which becomes (based on Part (i)) positive when $\bar{x} = \bar{x}_4$, and negative when $\bar{x} = \bar{x}_5$. $\square$

3.1 Local stability: the trivial equilibria

For the trivial equilibria $E_0 = (0, 0)$, $E_{\bar{x}_1} = E(\theta, 0)$ and $E_{\bar{x}_2} = E(k, 0)$, the Jacobian matrix in Eq. (3.1) implies

$$J(E_0) = \begin{bmatrix} e^{-r\theta} & 0 \\ 0 & e^{-d} \end{bmatrix}, \quad J(E_{\bar{x}_1}) = \begin{bmatrix} 1 + r\theta(1 - \frac{\theta}{k}) & \frac{-\theta}{\theta^2 + a} \\ 0 & G(\theta) \end{bmatrix} \quad \text{and} \\ J(E_{\bar{x}_2}) = \begin{bmatrix} 1 + r(\theta - k) & \frac{-k}{k^2 + a} \\ 0 & G(k) \end{bmatrix}.$$

Obviously, the eigenvalues in each case are located along the diagonal, which makes the stability obvious. Thus, we summarize the local stability in the following proposition.

Proposition 3.4. *Let $g(x) = \ln(G(x))$, and consider the three trivial equilibria with $\theta < k$, then we have*

[1] E_0 is always a sink.

[2] $E_{\bar{x}_1}$ is a source if $g(\theta) > 0$, a saddle if $g(\theta) < 0$, and non-hyperbolic if $g(\theta) = 0$.

[3] $E_{\bar{x}_2}$ is

- (i) a sink if $kr(\frac{\theta}{k} - 1) > -2$ and $g(k) < 0$;
- (ii) a source if $kr(\frac{\theta}{k} - 1) < -2$ and $g(k) > 0$;
- (iii) a saddle if either $kr(\frac{\theta}{k} - 1) > -2$ and $g(k) > 0$, or $kr(\frac{\theta}{k} - 1) < -2$ and $g(k) < 0$;
- (iv) a non-hyperbolic if $\frac{\mu k}{k^2 + a} = d$.

3.2 Local stability: one coexistence equilibrium exists

We need to deal with the two cases $\theta = \bar{x}_4 < \bar{x}_5 < k$ and $\theta < \bar{x}_4 < \bar{x}_5 = k$. Since Proposition 3.4 addresses the trivial equilibria, we need to focus on the nontrivial ones. Based on Proposition 3.1, we have to consider $E_5 = (\bar{x}_5, \bar{y}_5)$ when

$$\frac{\mu}{d} > 2\sqrt{a}, \quad \frac{\mu}{d} = \theta + \frac{a}{\theta} \quad \text{and} \quad \frac{\mu}{d} < k + \frac{a}{k}, \quad (3.5)$$

and $E_4 = (\bar{x}_4, \bar{y}_4)$ when

$$\frac{\mu}{d} > 2\sqrt{a}, \quad \frac{\mu}{d} < \theta + \frac{a}{\theta} \quad \text{and} \quad \frac{\mu}{d} = k + \frac{a}{k}. \quad (3.6)$$

Observe that Proposition 3.3 confines the options for each of E_5 and E_4 . Our next result addresses the local stability of these equilibria.

Theorem 3.5. *Consider the equilibria E_5 and E_4 as given in Eq. (3.5) and Eq. (3.6). Define*

$$\gamma_0(\theta; k, a, d) := dk\theta^5 - a(d+2)\theta^4 + 2ak(1-d)\theta^3 + 2a^2(d-1)\theta^2 + a^2k(d-6)\theta + (8-d)a^3.$$

Then, each of the following holds true:

- (i) E_5 is a source when $r\gamma_0(\theta, k; a, d) > 4k\theta^2(\theta^2 + a)$ and a saddle when the inequality is reversed.
- (ii) E_4 is a saddle when $r\gamma_0(k, \theta; a, d) > 4k^3(k^2 + a)$.
- (iii) E_4 is a sink when $r\gamma_0(k, \theta; a, d) < 4k^3(k^2 + a)$ and

$$k(k - \theta)[d(k^2 - a)^2 + ak^2 - 3a^2] < (k^2 - a)[d(k^2 - a)^2 - 4a^2].$$

- (iv) E_4 is a source when $r\gamma_0(k, \theta; a, d) < 4k^3(k^2 + a)$ and

$$k(k - \theta)[d(k^2 - a)^2 + ak^2 - 3a^2] > (k^2 - a)[d(k^2 - a)^2 - 4a^2].$$

Proof. The main idea here is to follow Lemma 3.2 and Fig. 2. In part (i), E_5 is a source when $T + D + 1 < 0$ and a saddle when $T + D + 1 > 0$. From the constraints we have on $\bar{x}_4$ and $\bar{x}_5$, we substitute

$$\bar{x}_5 = \frac{a}{\theta}, \quad \bar{y}_5 = r(\bar{x}_5^2 + a) \left(1 - \frac{\bar{x}_5}{k}\right) (\bar{x}_5 - \theta) \quad \text{and} \quad \mu = \frac{d}{\theta}(\theta^2 + a)$$

to obtain the result. Next, we verify Part (ii). From Proposition 3.3, we have $D > T - 1$ at E_4 . Thus, it remains to show that $D + T + 1 < 0$. From the constraints, we have

$$\bar{x}_4 = \frac{a}{k}, \quad \bar{y}_4 = r(\bar{x}_4^2 + a) \left(1 - \frac{\bar{x}_4}{k}\right) (\bar{x}_4 - \theta) \quad \text{and} \quad \mu = \frac{d}{k}(k^2 + a).$$

Substitute in $D + T + 1 < 0$ and simplify to obtain the result. To verify Part (iii), use the fact that $D > T - 1$ from Proposition 3.3, then from the condition $D + T + 1 > 0$ we obtain the inequality $r\gamma_0(k, \theta; a, d) < 4k^3(k^2 + a)$. Since $D - T + 1 > 0$ and $D + T + 1 > 0$ assure $D > -1$, we are left with the condition $D < 1$, which leads us to the second inequality. Finally, Part (iv) follows from $D + T + 1 > 0$ and $D > 1$. $\square$

Our next example shows specific cases that support the characterization given in Theorem 3.5.

Example 3.6. Consider the equilibria E_5 and E_4 as given in Eq. (3.5) and Eq. (3.6). Also, recall Part (ii) of Proposition 3.3.

(i) Fix $\mu = 4, a = 1$ and $d = \frac{8}{5}$, then $\theta = \bar{x}_4 = \frac{1}{2}$, $\bar{x}_5 = 2$ and $\bar{y}_5 = \frac{15}{2} \frac{r(k-2)}{k}$, $k > 2$. When fixing the parameters, keep in mind that we must have $k > \bar{x}_5 = \frac{\mu}{d} - \theta > \sqrt{a}$. From these fact and Part (i) of Theorem 3.5, we obtain E_5 is a saddle when $r(259 - 92k) < 50k$, and it is a source when the inequality is reversed. Both options are viable based on the values of $k > 2$ and $r > 0$.

(ii) Fix $d = 1, a = 14$ and $k = \bar{x}_5 = 4$. Then $\mu = \frac{15}{2}$, $\bar{x}_4 = \frac{7}{2}$, $\bar{y}_4 = \frac{105}{64} r(7 - 2\theta)$ and $0 < \theta < \frac{7}{2}$. Observe that we accommodated all conditions in (3.6). From these facts and Theorem 3.5, we obtain

- E_4 is a saddle when $\theta < \frac{1}{362r}(1057r - 960)$.
- E_4 is a sink when $\frac{1}{362r}(1057r - 960) < \theta < \frac{35}{12}$.
- E_4 is a source when $\theta > \max\{\frac{1}{362r}(1057r - 960), \frac{35}{12}\}$.

Figure 3: This figure shows a numerical simulation of the basin of attraction of the coexistence equilibrium $E_4(\bar{x}_4, \bar{y}_4) = (3.5, 4.9219)$, which has been indicated in red. The parameters are taken to be $d = 1$, $\mu = 7.5$, $a = 14$, $k = 4$, $r = 1$, and $\theta = 2$.

3.3 Local stability: two coexistence equilibria exist

Assume $\mu^2 > 4ad^2$, $\theta < \bar{x}_4$ and $\bar{x}_5 < k$. We focus on the local stability of $E_4 = (\bar{x}_4, \bar{y}_4)$ and $E_5 = (\bar{x}_5, \bar{y}_5)$. Based on Proposition 3.1, we need to proceed under the conditions

$$\frac{\mu}{d} > 2\sqrt{a} \quad \text{and} \quad \frac{\mu}{d} < \min \left\{ \theta + \frac{a}{\theta}, k + \frac{a}{k} \right\}. \quad (3.7)$$

As in the case of one coexistence equilibrium. Proposition 3.3 together with Lemma 3.2 show that the nontrivial equilibria merge at $a = \frac{\mu^2}{2d}$ as non-hyperbolic, and when they split, the stability of $E_5(\bar{x}_5, \bar{y}_5)$ lies below the line $D = T - 1$ in Figure 2, whereas the stability of $E_4(\bar{x}_4, \bar{y}_4)$ lies above the line $D = T - 1$. To get more insight into this matter, we depend on Figure 2 together with Lemma 3.2 and contemplate the signs of the expressions

$$\begin{aligned} D + T + 1 &= 2T + d\bar{y} \frac{(a - \bar{x}^2)}{(\bar{x}^2 + a)} \\ &= 4 + \frac{4r}{k} \bar{x} \left(\frac{k + \theta}{2} - \bar{x} \right) + d\bar{y} \left[\frac{4d}{\mu^2} + \frac{(a - \bar{x}^2)}{(\bar{x}^2 + a)} \right] \end{aligned} \quad (3.8)$$

and

$$D - 1 = \frac{2r}{k} \bar{x} \left(\frac{k + \theta}{2} - \bar{x} \right) + d\bar{y} \left[\frac{2d}{\mu^2} + \frac{(a - \bar{x}^2)}{(\bar{x}^2 + a)} \right]. \quad (3.9)$$

Although it is possible here to obtain an exact expression in terms of the parameters, we prefer to obtain an insight without writing explicit gigantic expressions. The signs of the expressions $D + T + 1$ and $D - 1$ are determined by whether $\bar{x} = \bar{x}_4$ or $\bar{x} = \bar{x}_5$, and how far is $\bar{x}$ from the

average of k and θ . In particular, when $\bar{x} = \bar{x}_4$, having $\bar{x}_4$ close to the Allee parameter θ makes $E_4(\bar{x}_4, \bar{y}_4)$ a source. On the other hand, having $\bar{x}_4$ “close” to the carrying capacity of the prey can make $E_4(\bar{x}_4, \bar{y}_4)$ a sink, while having $\bar{x}_4$ “too close” to the carrying capacity k of the prey can make $E_4(\bar{x}_4, \bar{y}_4)$ a saddle. When $\bar{x} = \bar{x}_5$, having $\bar{x}_5$ “close” to the Allee parameter θ makes $E_5(\bar{x}_5, \bar{y}_5)$ a source, while having $\bar{x}_5$ “close” to the carrying capacity k of the prey can make $E_5(\bar{x}_5, \bar{y}_5)$ a saddle. Our next example illustrates these scenarios:

Example 3.7. Consider $d = 1$, $\mu = 17$, $a = 72$ and $k = 10$. We obtain

$$\bar{x}_4 = 8, \bar{y}_4 = \frac{136}{5}r(8 - \theta), \bar{x}_5 = 9 \quad \text{and} \quad \bar{y}_5 = \frac{153}{10}r(9 - \theta).$$

At $E_4(\bar{x}_4, \bar{y}_4)$, we have

$$T(\theta) = \frac{52}{85}r\theta + 2 - \frac{56}{17}r \quad \text{and} \quad D(\theta) = \frac{3}{5}r\theta + 1 - \frac{16}{5}r.$$

For $r > \frac{85}{138}$ and “small” values of θ , $E_4 := E_4(\bar{x}_4, \bar{y}_4)$ is a saddle with $\lambda_1 < -1$ and $-1 < \lambda_2 < 1$. As we increase θ , E_4 becomes nonhyperbolic at $\theta = \frac{1}{103}(1 - \frac{340}{r})$, which only takes place when $r > 340$, then followed by a sink with real eigenvalues, then a sink with complex eigenvalues at

$$\theta = \frac{1}{1352r} (7280r - 85 - 17\sqrt{2080r + 25}),$$

then nonhyperbolic again at $\theta = \frac{16}{3}$ followed by a source with complex eigenvalues, then nonhyperbolic again at

$$\theta = \frac{1}{1352r} (7280r - 85 + 17\sqrt{2080r + 25})$$

followed by a source with real eigenvalues. By plotting the curve $(T(\theta), D(\theta))$, $0 \leq \theta \leq 8$, Fig. 4 can be used to illustrate these scenarios. On the other hand, at $E_5(\bar{x}_5, \bar{y}_5)$, we have

$$T(\theta) = \frac{27}{34}r\theta + 2 - \frac{531}{85}r \quad \text{and} \quad D(\theta) = \frac{4}{5}r\theta + 1 - \frac{63}{10}r.$$

As the previous example shows, we have a wide range of scenarios. Therefore, we proceed by limiting our attention to the particular case

$$k = \theta + 10, \quad \mu = d(2\theta + 13) \quad a = (\theta + 6)(\theta + 7). \quad (3.10)$$

The motivation for this restrictions stems from the interesting values of the equilibria, namely

$$\begin{aligned} \bar{x}_4 &= \frac{\theta + k}{2} + 1 = \theta + 6, \quad \bar{y}_4 = \frac{24r(\theta + 6)(2\theta + 13)}{\theta + 10} \quad \text{and} \\ \bar{x}_5 &= \frac{\theta + k}{2} + 2 = \theta + 7, \quad \bar{y}_5 = \frac{21r(\theta + 7)(2\theta + 13)}{\theta + 10}. \end{aligned}$$

Observe that this restriction accommodates the conditions in (3.7).

Next, we give the following two results, which are useful in the bifurcation analysis.

Proposition 3.8. Assume the conditions in (3.7) are satisfied. Each of the following holds true:

- (i) If $2\bar{x}_4 < \theta + k$, then E_4 is a source.

Figure 4: Figure (a) shows the plot of the eigenvalues captured at $r = 2$, while Figure (b) shows the stability of $E_4(\bar{x}_4, \bar{y}_4)$ in the (r, θ) -plane.

Figure 5: This figure shows a numerical simulation of the basin of attraction of the equilibrium point $E_4(\bar{x}_4, \bar{y}_4) = (8, 136)$, which has been indicated by the red shaded circle. Here $d = 1$, $\mu = 17$, $a = 72$, $k = 10$, $r = 1$, and $\theta = 3$.

(ii) Suppose that $\frac{1}{3} \left(\theta + k + \sqrt{(k - \theta)^2 + k\theta} \right) < \frac{\mu}{2d}$. There exists $\epsilon > 0, r_0 > 0$ and a , such that $\frac{1}{2}(k + \theta) < \sqrt{a} < k$ and E_4 is a saddle whenever $r \geq r_0$ and $\bar{x}_4 \in (\sqrt{a} - \epsilon, \sqrt{a})$.

Proof. (i) Suppose that $\bar{x}_4 \leq \frac{1}{2}(k + \theta)$. From Proposition 3.3, we always have $D - T + 1 > 0$ at E_4 . Also, from equation (3.8) and (3.9), we obtain $D > 1$, which makes E_4 a source based on Lemma

3.2. To prove Part (ii), recall that we have

$$\theta < \bar{x}_4 < \sqrt{a} < \bar{x}_5 < k \quad \text{and} \quad \bar{x}_4 = \sqrt{a} \quad \text{at} \quad a = \frac{\mu}{2d}.$$

From Proposition 3.3, we always have $D - T + 1 > 0$ at E_4 , and from Eq. (3.9), we have

$$D + T + 1 = 4 + \frac{4r}{k} \bar{x}_4 \left(\frac{k + \theta}{2} - \bar{x}_4 \right) + d\bar{y}_4 \left[\frac{4d}{\mu^2} + \frac{(a - \bar{x}_4^2)}{(\bar{x}_4^2 + a)^2} \right].$$

At $\bar{x}_4 = \sqrt{a}$, we obtain

$$D + T + 1 = 4 - \frac{2}{k} \left[3 \left(\frac{\mu}{2d} \right)^2 - 2(k + \theta) \frac{\mu}{2d} + k\theta \right] r. \quad (3.11)$$

Now, to have the coefficient of r negative and keep the inequalities in (2.7) satisfied, we need

$$\frac{1}{3} \left(\theta + k + \sqrt{(k - \theta)^2 + k\theta} \right) < \frac{\mu}{2d} < k,$$

which has a feasible region all the time. Finally,

$$r \geq \frac{2k}{3 \left(\frac{\mu}{2d} \right)^2 - 2(k + \theta) \frac{\mu}{2d} + k\theta} =: r_0 > 0$$

makes $D + T + 1 < 0$, which completes the proof. $\square$

In Part (ii) of Proposition 3.8, we confined $\bar{x}_4$ to make E_4 a saddle, and from Part (i), θ can change E_4 into a source. In the next result, we force any change from a source to a saddle in E_4 to go through the sink case.

Lemma 3.9. *If $d \leq \frac{16}{rk}$, then a change in E_4 from a source to a saddle (or visa-versa) goes through the sink case.*

Proof. To guarantee that a transition from a saddle to a source goes through the sink case, we focus on the expression

$$D - T + 1 = \frac{d\bar{y}_4(a - \bar{x}_4^2)}{(a + \bar{x}_4^2)^2}.$$

The right hand side is obviously positive; however, for the line to pass through the sink region, we need the right hand side to be less than 4. Thus, substitute the value of $\bar{y}_4$ and use the given condition on d to obtain

$$\begin{aligned} \frac{dr}{k} \frac{(k - \bar{x}_4)(\bar{x}_4 - \theta)(a - \bar{x}_4^2)}{(a + \bar{x}_4^2)} &\leq \frac{16}{k^2} (k - \bar{x}_4)(\bar{x}_4 - \theta) \\ &\leq \frac{16}{k^2} \max\{(k - x)(x - \theta) : \theta < x < k\} \\ &= \frac{4}{k^2} (k - \theta)^2 < 4. \end{aligned}$$

This validates our statement. $\square$

4 Bifurcation analysis

In this section, we assume the existence of two coexistence equilibria $\theta < \bar{x}_4 < \bar{x}_5 < k$ as in Subsection 3.3. Thus, we again stress the needed conditions in (3.7). From the stability analysis provided in this case, the nontrivial equilibrium E_4 is expected to go through period-doubling (flip) bifurcation when the parameter values force E_4 to exit the sink zone into the saddle zone (or period-halving bifurcation if the direction is reversed). Also, E_4 is expected to go through a Neimark-Sacker (Hopf) bifurcation when the parameter values force E_4 to exit the sink zone into the source zone through the line segment $D = 1$. See Fig. 2 for elaboration. In this section, we consider the Allee parameter θ as the bifurcation parameter, then prove the period-doubling and Neimark-Sacker bifurcations of the nontrivial equilibrium point E_4 within certain choices of the parameters. As can be observed from Proposition 3.8 and Lemma 3.9, we can have a change from a saddle to a sink, or a sink to a source, for a wide choice of the parameters. Since our main interest is on the effect of r and θ , we avoid the inconvenience of having awkward expressions by confining some of the parameters. Thus, we proceed our bifurcation analysis with the following restrictions

$$d := \frac{16}{rk}, \quad \mu := \frac{32}{k}, \quad a := \frac{\mu^2}{4d^2} - 1 \quad \text{and} \quad k := r + 2. \quad (4.1)$$

Under these restrictions, we have $\bar{x}_4 = r - 1$ and $\bar{x}_5 = r + 1 < k$. To have two coexistence equilibria, we accommodate the conditions in 3.7, and therefore, restrict the Allee parameter $\theta < r - 1$.

Theorem 4.1. *Let $\theta < r - 1$, and consider d, μ, a and k as restricted in equations (4.1), then each of the following holds true:*

- (i) *There exists $r_0 > 1$ and $\theta = \theta_0(r)$, $r > r_0$ such that the Jacobian matrix at E_4 has $\lambda_1 = -1$ and $|\lambda_2| < 1$. More specifically,*

$$\theta_0(r) := r - 4 - \frac{11r^3 + 17r^2 - 10r + 72}{r^4 - 2r^3 - 5r^2 + 6r - 24}.$$

- (ii) *There exists $r_0 > 1$ and $\theta = \theta_1(r)$, $r > r_0$, such that the Jacobian matrix at E_4 has both eigenvalues non-real with modulus equal one. More specifically,*

$$\theta_1(r) = r - 4 - \frac{9r^3 + 9r^2 - 18r + 144}{r^4 - 2r^3 - 5r^2 + 6r - 48}.$$

Furthermore, θ_0 and θ_1 satisfy $0 < \theta_0 < \theta_1 < r - 1$.

Proof. Under the confined values of d, μ, a and k , the trace-determinant values at E_4 give us

$$D - T + 1 = \frac{48(r - 1 - \theta)}{r(r + 2)^2}. \quad (4.2)$$

Since $\theta < r - 1$, the right hand side is positive. Also, from Lemma 3.9, the right hand side is smaller than 4. Now, $D + T + 1 = 0$ implies

$$\theta = \theta_0(r) := r - 4 - \frac{11r^3 + 17r^2 - 10r + 72}{r^4 - 2r^3 - 5r^2 + 6r - 24}, \quad (4.3)$$

and $D - 1 = 0$ implies

$$\begin{aligned}\theta = \theta_1(r) &:= (r - 1) \left(1 - \frac{3r^2(r + 2)}{r^4 - 2r^3 - 5r^2 + 6r - 48} \right) \\ &= r - 4 - \frac{9r^3 + 9r^2 - 18r + 144}{r^4 - 2r^3 - 5r^2 + 6r - 48}.\end{aligned}\quad (4.4)$$

Now, it is a matter of elementary computations to conclude parts (i) and (ii), and the consequent fact $0 < \theta_0 < \theta_1 < r - 1$. $\square$

4.1 Neimark-Sacker bifurcation

We proceed in this section to prove the Neimark-Sacker bifurcation of the equilibrium point E_4 . Again, we avoid awkward expressions by considering the restrictions in equations (4.1). From Theorem 4.1, the determinant and trace of the Jacobian in Eq. (3.3) satisfy $D(\theta) = 1$ and $-2 < T(\theta) < 2$ at $\theta = \theta_1$. Let θ vary in a sufficiently small neighborhood of θ_1 , and write the eigenvalues of J at E_4 as

$$\lambda_{1,2}(\theta) = \frac{1}{2}T(\theta) \pm \frac{1}{2}\sqrt{T^2(\theta) - 4D(\theta)}, \quad \lambda_{1,2}(\theta_1) = \frac{1}{2}T(\theta_1) \pm i\frac{1}{2}\sqrt{4 - T^2(\theta_1)}, \quad (4.5)$$

we give the following result.

Theorem 4.2. *Consider the assumptions of Theorem 4.1, then the model in Eq. (2.1) goes through a Neimark-Sacker bifurcation at $\theta = \theta_1$.*

Proof. Theorem 4.1 assures the existence of θ_1 at which the eigenvalues are complex with modulus one. It remains to show that

$$\frac{d}{d\theta} |\lambda_{1,2}(\theta_1)| \neq 0 \quad \text{and} \quad \lambda_{1,2}^m(\theta_1) \neq 1 \quad \text{for} \quad m = 1, 2, 3, 4.$$

Since

$$|\lambda_{1,2}(\theta)| = \sqrt{D(\theta)}, \quad \text{we obtain} \quad \frac{d}{d\theta} |\lambda_{1,2}(\theta)| = \frac{D'(\theta)}{2\sqrt{D(\theta)}},$$

and consequently

$$\begin{aligned}\frac{d}{d\theta} |\lambda_{1,2}(\theta_h)| &= \frac{1}{2}D'(\theta_1) \\ &= \frac{r((3-d)\bar{x}_4^3 + k(d-2)\bar{x}_4^2 + a(d+1)\bar{x}_4 - adk)}{(\bar{x}_4^2 + a)k} \\ &= r - 6 + \frac{3(5r^2 + 10r - 16)}{r(r+2)^2},\end{aligned}$$

Rather than showing the right hand side cannot be zero directly, we assume $\frac{d}{d\theta} |\lambda_{1,2}(\theta_1)| = 0$, and use the equation $D(\theta_1) = 1$ as obtained in Eq. (4.2) to eliminate r using the resultant. In this case, we obtain $429981696 = 0$, which is a contradiction. Therefore, $\frac{d}{d\theta} |\lambda_{1,2}(\theta_h)|$ cannot be zero. Next, we solve $\lambda_{1,2}^m(\theta_1) = 1$. When $m = 1$, $\lambda_{1,2}(\theta_1) = 1$ if and only if $T = 2$, which is not possible from Eq. (4.2) and Theorem 4.1. When $m = 2$, $\lambda_{1,2}^2(\theta_1) = 1$ if and only if $T = 2$ or -2 . $T = 2$ is not possible by our previous argument, and $T = -2$ is not possible from Lemma 3.9. When $m = 3$,

$\lambda_{1,2}^3(\theta_1) = 1$ if and only if $T = 2, -1, -1$. When $m = 4$, $\lambda_{1,2}^4(\theta_1) = 1$ if and only if $T = 2, -2, 0, 0$. Thus, it remains to show that $(T, D) = (0, 1), (-1, 1)$ are both not possible. For $(T, D) = (0, 1)$, substitute and eliminate θ_1 to obtain

$$2x^5 + 10x^4 + 2x^3 - 186x^2 - 276x - 288 = 0, \quad \text{where } x = r - 1.$$

which has only one real solution between $r = 4$ and $r = 5$. Also, for $(T, D) = (-1, 1)$, we substitute and eliminate θ_1 to obtain

$$3x^5 + 15x^4 + 3x^3 - 207x^2 - 342x - 432 = 0, \quad \text{where } x = r - 1.$$

□

Now, the center manifold reduction technique can be used to show the existence and stability of the Neimark-Sacker invariant curve [29; 30]. We avoid the lengthy computations and give the following numerical example based on the obtained theoretical results.

Example 4.3. Consider $r = 8$, $k = r + 2 = 10$, $d = \frac{16}{rk} = \frac{1}{5}$ and $\mu = \frac{32}{k} = \frac{16}{5}$. In this case, we have $\bar{x}_4 = r - 1 = 7$, $\bar{y}_4 = \frac{1}{5}(9408 - 1344\theta)$ and $\bar{x}_5 = r + 1 = 8$, $\bar{y}_5 = \frac{1}{5}(5184 - 576\theta)$. From Eq. (4.4), we have $\theta_1 = \frac{91}{43} \approx 2.116279$. At this value of θ , we have $E_4 = (7, \frac{56448}{43}) \approx (7, 1312.744)$ and $E_5 = (8, \frac{170496}{215}) \approx (8, 793.005)$. The eigenvalues at E_4 and E_5 are given by $\frac{367}{430} \pm \frac{3}{430}\sqrt{5579}i$, $-\frac{3673}{215} \pm \frac{2}{215}\sqrt{3780727}$, respectively. Fig. 6 shows a numerical simulation of the Neimark-Sacker invariant curve at θ near θ_1 .

Figure 6: This figure shows the existence of a Neimark-Sacker bifurcation at θ near θ_1 . Namely $\theta = 2.12$. The other parameters are taken as in Example 4.3. The black curve indicates part of the nullcline of the prey equation.

4.2 Period-doubling bifurcation

Recall the restrictions we put on our parameters as given in inequalities (4.1), and consider the equilibrium point E_4 . We focus on the period-doubling (or flip) bifurcation of our Model (2.1) at E_4 . In Theorem 4.1, we provided sufficient conditions to guarantee the existence of $\theta = \theta_0$ at which $D+T+1 = 0$ and $-2 < T < 0$. This is the necessary assumption to start with for a period-doubling bifurcation. Now, we proceed based on the center manifold reduction and the corresponding critical normal form to test the super-criticality of the newly born 2-cycle. Now, we shift the equilibrium point E_4 into the origin by letting $(u_n, v_n) = (x_n - \bar{x}_4, y_n - \bar{y}_4)$. Thus, we have

$$\begin{cases} u_{n+1} = F_1(u_n, v_n) := (u_n + \bar{x}_4)F(u_n + \bar{x}_4, v_n + \bar{y}_4) - \bar{x}_4 \\ v_{n+1} = F_2(u_n, v_n) := (v_n + \bar{y}_4)G(u_n + \bar{x}_4) - \bar{y}_4. \end{cases} \quad (4.6)$$

Write $X := [u, v]^t$ and $\bar{X} := (\bar{u}, \bar{v}) = (0, 0)$, then use Taylor expansion at $\bar{X}$ (up to terms of order 3). For writing conveniences, we follow the notations of Kuznetsov [29] (Chapter 4). Let q be the eigenvector that belongs to the eigenvalue $\lambda_1 = -1$, and let p be the adjoint eigenvector, i.e., $J(E_4)q = \lambda_1 q$ and $J^T(E_4)p = \lambda_1 p$, respectively. Normalize p with respect to q such that $\langle p, q \rangle = 1$ and $\langle \cdot, \cdot \rangle$ is the standard inner product on $\mathbb{R}^2$. The normal form of our system becomes

$$\xi \mapsto -\xi + \frac{1}{6}c\xi^3 + O(\xi^4),$$

where

$$\begin{aligned} c &:= \langle p, C(q, q, q) - 3B(q, (J - I_2)^{-1}B(q, q)) \rangle \\ B(q, q) &= [B_1(q_1, q_2), B_2(q_1, q_2)]^T \\ C(q, q, q) &= [C_1(q_1, q_2), C_2(q_1, q_2)]^T \end{aligned}$$

and

$$\begin{aligned} B_i(q_1, q_2) &= \frac{\partial^2 F_i(0, 0)}{\partial x^2} q_1^2 + 2 \frac{\partial^2 F_i(0, 0)}{\partial x y} q_1 q_2 + \frac{\partial^2 F_i(0, 0)}{\partial y^2} q_2^2 \\ C_i(q_1, q_2) &= \frac{\partial^3 F_i(0, 0)}{\partial x^3} q_1^3 + 3 \frac{\partial^3 F_i(0, 0)}{\partial x^2 \partial y} q_1^2 q_2 + 3 \frac{\partial^3 F_i(0, 0)}{\partial y^2 \partial x} q_1 q_2^2 + \frac{\partial^3 F_i(0, 0)}{\partial y^3} q_2^3. \end{aligned}$$

Now, we do the computational part. We start by the Jacobian matrix at E_4 and $\theta = \theta_0$. Use the fact that $\lambda_1 = -1$ and write $\lambda_2 = \lambda$, we obtain

$$J(E_4) = \begin{bmatrix} \lambda - 2 & -\frac{d}{\mu} \\ \frac{2\mu}{d}(1 - \lambda) & 1 \end{bmatrix}. \quad (4.7)$$

The eigenvectors p and q are given by

$$\begin{aligned} q &= \begin{bmatrix} q_1 & q_2 \end{bmatrix}^T = \begin{bmatrix} \frac{d}{\mu} & \lambda - 1 \end{bmatrix}^T = \begin{bmatrix} \frac{1}{2r} & \lambda - 1 \end{bmatrix}^T \\ p &= \begin{bmatrix} p_1 & p_2 \end{bmatrix}^T = \frac{1}{\lambda + 1} \begin{bmatrix} 2\frac{\mu}{d} & 1 \end{bmatrix}^T = \frac{1}{\lambda + 1} \begin{bmatrix} 4r & 1 \end{bmatrix}^T. \end{aligned}$$

We evaluate $B(q, q)$ and $C(q, q, q)$ to obtain

$$B(q, q) = \begin{bmatrix} B_1(q_1, q_2) \\ B_2(q_1, q_2) \end{bmatrix} = \begin{bmatrix} -\frac{(2r^7-6r^6+3r^5+19r^4-84r^3+204r^2-96r+192)}{4r^3(r+2)(r^4-2r^3-5r^2+6r-24)} \\ -\frac{24(3r^2-r+4)}{r^2(r^4-2r^3-5r^2+6r-24)} \end{bmatrix}$$

and

$$C(q, q, q) = \begin{bmatrix} C_1(q_1, q_2) \\ C_2(q_1, q_2) \end{bmatrix} = \begin{bmatrix} \frac{(6r^9-24r^8+31r^7-6r^6-327r^5+1154r^4-1830r^3+1860r^2-1728r+576)}{8(r^5(r-1)^2(r+2)(r^4-2r^3-5r^2+6r-24))} \\ \frac{(12(3r^2-r+4))(3r^7+9r^6-6r^5-54r^4-72r^3+24r^2+96r-128)}{(r^6(r-1)^2(r+2)^3(r^4-2r^3-5r^2+6r-24))} \end{bmatrix}$$

Next, we evaluate $B(q, (J - I_2)^{-1}B(q, q))$. We have

$$(J - I_2)^{-1}B(q, q) = \begin{bmatrix} Q_1 \\ Q_2 \end{bmatrix} = \begin{bmatrix} -\frac{(r+2)}{4r^3} \\ \frac{(2r^7-4r^6+7r^5+r^4-56r^3+284r^2-192r+192)}{2r^2(r^4-2r^3-5r^2+6r-24)(r+2)} \end{bmatrix}.$$

Therefore,

$$B(q, (J - I_2)^{-1}B(q, q)) = \begin{bmatrix} \frac{(2r^9-17r^7+52r^6-76r^5-175r^4+732r^3-956r^2+624r-384)}{8(r^5(r-1)(r+2)(r^4-2r^3-5r^2+6r-24))} \\ \frac{(4(11r^6+38r^5+58r^4+7r^3-116r^2+140r-336))}{(r^4(r-1)(r+2)^2(r^4-2r^3-5r^2+6r-24))} \end{bmatrix}.$$

Finally, we evaluate $C(q, q, q) - 3B(q, (J - I_2)^{-1}B(q, q))$ and do the inner product to obtain

$$c = \frac{R_2(r) - R_1(r)}{4r^6(r^5 - 9r^3 - 40r^2 - 96)(r + 2)^2(r - 1)^2}, \quad (4.8)$$

where

$$\begin{aligned} R_1(r) &:= 6r^{14} + 12r^{13} + 20r^{11} + 482r^{10} + 182r^9 + 1375r^8 + 3250r^7 + 1992r^5 + 9504r^4 + 27648r^2 + 12288 \\ R_2(r) &:= 51r^{12} + 6308r^6 + 17088r^3 + 12288r. \end{aligned}$$

Based on the above facts, we summarize in the following result.

Theorem 4.4. *Consider d, μ, a and k as restricted in equations (4.1), then the model in Eq. (2.1) undergoes a period doubling bifurcation at the equilibrium point E_4 when θ varies in a small neighborhood of θ_0 . Furthermore, the stability of the newly born 2-cycle depends on the sign of c in Eq. (4.8).*

5 Conclusion

Dynamical complexities are common characteristic in different type of population models. Different types of biological phenomena can impact dynamical behaviors of a model. The stability analysis of predator-prey system has great importance in understanding its biological relevance and may be helpful in conservation management. Allee effect is an important ecological phenomenon and could have stabilizing or destabilizing effect on population dynamics in general. In this paper,

we considered the strong Allee effect for the prey growth function in a predator-prey system with non-monotonic functional response. We have performed stability analysis and provided sufficient conditions for the period-doubling and the Neimark-Sacker bifurcations by taking Allee parameter as bifurcation parameter. The model without Allee effect studied by Hu et al. [31] and Wang and Li [32], and they observed chaotic dynamics in the system. We found that Allee effect has a stabilizing effect on the system for certain values of the parameters. It is observed that replacing the logistic growth function with Allee type function changes the dynamical behavior of the system. So, Allee effect may be used as a stabilizing factor in management policy of species conservation.

References

- [1] R. M. May. Simple mathematical models with very complicated dynamics. *Nature*, 261:459–467, 1976.
- [2] Robert M. May. Biological populations with nonoverlapping generations: Stable points, stable cycles, and chaos. *Science*, 186(4164):645–647, 1974.
- [3] H. Al-Kahby, F. Dannan, and S. Elaydi. Non-standard discretization methods for some biological models. In *Applications of nonstandard finite difference schemes (Atlanta, GA, 1999)*, pages 155–180. World Sci. Publ., River Edge, NJ, 2000.
- [4] A. Hastings and T. Powell. Chaos in three-species food chain. *Ecology*, 72:896–903, 1991.
- [5] AJ Lotka. Elements of physical biology, williams and wilkins, baltimore. 1925.
- [6] Vito Volterra. Fluctuations in the abundance of a species considered mathematically. 118:558–560, 1926.
- [7] W. C. Allee. *Animal Aggregations: A study in General Sociology*. University of Chicago Press, USA, 1931.
- [8] F. Courchamp, L. Berec, and J. Gascoigne. *Allee effects in ecology and conservation*. Oxford University Press, Oxford, 2008.
- [9] Philip A Stephens and William J Sutherland. Consequences of the Allee effect for behaviour, ecology and conservation. *Trends in ecology & evolution*, 14(10):401–405, 1999.
- [10] Franck Courchamp, Tim Clutton-Brock, and Bryan Grenfell. Inverse density dependence and the allee effect. *Trends in ecology & evolution*, 14:405–410, 1999.
- [11] J. B. Ferdy, F. Austerlitz, J. Moret, P. H. Gouyon, and B. Godelle. Pollinator-induced density dependence in deceptive species. *Oikos*, 87:549–560, 1999.
- [12] M. J Groom. Allee effects limit population viability of an annual plant. *The American Naturalist*, 151(6):487–496, 1998.

- [13] M. Kuussaari, I. Saccheri, M. Camara, and I. Hanski. Allee effect and population dynamics in the glanville fritillary butterfly. *Oikos*, 82:384–392, 1998.
- [14] A. W Stoner and M. Ray-Culp. Evidence for Allee effects in an over-harvested marine gastropod: density-dependent mating and egg production. *Marine Ecology Progress Series*, 202:297–302, 2000.
- [15] F. Courchamp, B.T. Grenfell, and T.H. Clutton-Brock. Impact of natural enemies on obligately cooperative breeders. *Oikos*, 91:311–322, 2000.
- [16] P. Amarasekare. Interactions between local dynamics and dispersal: Insights from single species models. *Theoretical Population Biology*, 53(1):44 – 59, 1998.
- [17] J. M Drake. Allee effects and the risk of biological invasion. *Risk Analysis*, 24:795–802, 2004.
- [18] J. Shi and R. Shivaji. Persistence in reaction diffusion models with weak Allee effect. *Journal of Mathematical Biology*, 52:807–829, 2006.
- [19] C. M. Taylor and A. Hastings. Allee effects in biological invasions. *Ecology Letters*, 8:895–908, 2005.
- [20] C. Celik and O. Duman. Allee effect in a discrete-time predator–prey system. *Chaos, Solitons & Fractals*, 40:1956–1962, 2009.
- [21] W. X. Wang, Y. B. Zhang, and C.Z. Liu. Analysis of a discrete-time predator–prey system with Allee effect. *Ecological Complexity*, 8:81–85, 2011.
- [22] Saheb Pal, Sourav Kumar Sasmal, and Nikhil Pal. Chaos control in a discrete-time predator–prey model with weak Allee effect. *International Journal of Biomathematics*, 11(07):1850089, 2018.
- [23] Saheb Pal, Nikhil Pal, and Joydev Chattopadhyay. Hunting cooperation in a discrete-time predator–prey system. *International Journal of Bifurcation and Chaos*, 28(07):1850083, 2018.
- [24] W Sokol and JA Howell. Kinetics of phenol oxidation by washed cells. *Biotechnology and Bioengineering*, 23(9):2039–2049, 1981.
- [25] Dongmei Xiao and Shigui Ruan. Global analysis in a predator–prey system with nonmonotonic functional response. *SIAM Journal on Applied Mathematics*, 61(4):1445–1472, 2001.
- [26] L. Dai. *Nonlinear dynamics of piecewise constant systems and implementation of piecewise constant arguments*. World Scientific, 2008.
- [27] Sebastian J Schreiber. Allee effects, extinctions, and chaotic transients in simple population models. *Theoretical population biology*, 64(2):201–209, 2003.
- [28] Saber N Elaydi and Robert J Sacker. Population models with allee effect: a new model. *Journal of Biological Dynamics*, 4(4):397–408, 2010.

- [29] Yuri A. Kuznetsov. *Elements of applied bifurcation theory*, volume 112 of *Applied Mathematical Sciences*. Springer-Verlag, New York, third edition, 2004.
- [30] J. Guckenheimer and P. J Holmes. *Nonlinear oscillations, dynamical systems, and bifurcations of vector fields*, volume 42. Springer Science & Business Media, 2013.
- [31] Z. Hu, Z. Teng, and L. Zhang. Stability and bifurcation analysis of a discrete predator-prey model with non-monotonic functional response. *Nonlinear Analysis: Real World Applications*, 12:2356–2377, 2011.
- [32] C. Wang and X. Li. Further investigations into the stability and bifurcation of a discrete predator–prey model. *Journal of Mathematical Analysis and Applications*, 422:920–939, 2015.